


Mark Scheme (Results) Summer 2008

GCE

GCE Mathematics (6674/01)




**June 2008
Further Pure FP1
Mark Scheme**

Question number	Scheme	Marks
1.	<p>(a) 4</p> <p>(b) $(x-4)(x^2+4x+16)$</p> $x = \frac{-4 \pm \sqrt{16-64}}{2}, \quad x = -2 \pm 2\sqrt{3}i \quad (\text{or equiv. surd for } 2\sqrt{3})$ <p>(c)  <p style="margin-left: 100px;">Root on +ve real axis, one other in correct quad.</p> <p style="margin-left: 100px;">Third root in conjugate complex position</p> </p>	<p>B1 (1)</p> <p>M1 A1</p> <p>M1, A1 (4)</p> <p>B1</p> <p>B1ft (2)</p> <p style="text-align: right;">7</p>
	<p>M1 in part (b) needs $(x-4)$ times quadratic $(x^2+ax+..)$ or times (x^2+16)</p> <p>M1 needs solution of three term quadratic</p> <p>So (x^2+16) special case, results in B1M1A0M0A0B0B1 possibly</p> <p>Alternative scheme for (b)</p> <p>$(a+ib)^3 = 64$, so $a^3 + 3a^2ib + 3a(ib)^2 + (ib)^3 = 64$ and equate real, imaginary parts</p> <p style="margin-left: 40px;">so $a^3 - 3ab^2 = 64$ and $3a^2b - b^3 = 0$</p> <p style="margin-left: 40px;">Solve to obtain $a = -2, b = \sqrt{12}$</p> <p>Alternative ii</p> <p>$(x-4)(x-a-ib)(x-a+ib) = 0$ expand and compare coefficients</p> <p>two of the equations $-2a-4=0, 8a+a^2+b^2=0, 4(a^2+b^2)=64$</p> <p style="margin-left: 40px;">Solve to obtain $a = -2, b = \sqrt{12}$</p> <p>(c) Allow vectors, line segments or points in Argand diagram.</p> <p>Extra points plotted in part (c) – lose last B mark</p> <p>Part (c) answers are independent of part (b)</p>	<p>M1</p> <p>A1</p> <p>M1A1</p> <p>M1</p> <p>A1</p> <p>M1A1</p>

Question number	Scheme	Marks
2.	<p>(a) $f(1.6) = \dots$ $f(1.7) = \dots$ (Evaluate both)</p> <p>0.08... (or 0.09), -0.3... One +ve, one -ve or Sign change, \therefore root</p> <p>(b) $f'(x) = -4\sin x - e^{-x}$</p> <p>$1.6 - \frac{f(1.6)}{f'(1.6)}$</p> <p>$= 1.6 - \frac{4\cos 1.6 + e^{-1.6}}{(-4\sin 1.6 - e^{-1.6})}$ $\left(= 1.6 - \frac{0.085\dots}{-4.2\dots} \right)$</p> <p>$= 1.62$</p>	<p>M1</p> <p>A1 (2)</p> <p>B1</p> <p>M1</p> <p>A1</p> <p>A1 (4)</p> <p>6</p>
	<p>(a) Any errors seen in evaluation of $f(1.6)$ or $f(1.7)$ lose A mark so -0.32 is A0 Values are 0.0851 and -0.3327 Need concluding statement also.</p> <p>(b) B1 may be awarded if seen in N-R as $-4\sin 1.6 - e^{-1.6}$ or as -4.2 M1 for statement of Newton Raphson (sign error in rule results in M 0) First A1 may be implied by correct work previously followed by correct answer Do not accept 1.620 for final A1. It must be given and correct to 3sf. 1.62 may follow incorrect work and is A0 No working at all in part (b) is zero marks.</p>	

Question number	Scheme	Marks
3.	<p>(a) $z = \frac{(a+2i)(a+i)}{(a-i)(a+i)} = \frac{a^2 + 3ai - 2}{a^2 + 1}$</p> <p>$\frac{a^2 - 2}{a^2 + 1} = \frac{1}{2}, \quad 2a^2 - 4 = a^2 + 1 \quad a = \sqrt{5} \quad (\text{presence of } -\sqrt{5} \text{ also is A0})$</p> <p>(b) Evaluating their “$\frac{3a}{a^2 + 1}$”, or “$3a$” $\left(\frac{\sqrt{5}}{2} \text{ or } 3\sqrt{5}\right)$ (ft errors in part a)</p> <p>$\tan \theta = \frac{3a}{a^2 - 2} (= \frac{3\sqrt{5}}{3}), \arg z = 1.15 \quad (\text{accept answers which round to 1.15})$</p>	<p>M1 A1</p> <p>M1, A1 (4)</p> <p>B1ft</p> <p>M1, A1 (3)</p> <p>7</p>
	<p>(b) B mark is treated here as a method mark</p> <p>The M1 is for $\tan(\arg z) = \text{Imaginary part} / \text{real part}$</p> <p>answer in degrees is A0</p> <p><u>Alternative method:</u></p> <p>(a) $\left(\frac{1}{2} + iy\right)(a - i) = a + 2i \Rightarrow \frac{1}{2}a + y = a \text{ and } ay - \frac{1}{2} = 2$</p> <p>$y = \frac{1}{2}a \text{ and } ay = \frac{5}{2} \Rightarrow \frac{1}{2}a^2 = \frac{5}{2} \Rightarrow a = \sqrt{5}$</p> <p>(b) $y = \frac{\sqrt{5}}{2}$ (May be seen in part (a))</p> <p>$\tan \theta = \sqrt{5} \quad \arg z = 1.15$</p> <p><u>Further Alternative method in (b)</u></p> <p>Use $\arg(a + 2i) - \arg(a - i)$</p> <p>$= 0.7297 - (-0.4205) = 1.15$</p>	<p>M1 A1</p> <p>M1 A1 (4)</p> <p>B1ft</p> <p>M1 A1 (3)</p> <p>B1</p> <p>M1A1 (3)</p>

Question number	Scheme	Marks
4.	<p>(a) $m^2 + 4m + 3 = 0$ $m = -1, m = -3$</p> <p>C.F. $(x =) Ae^{-t} + Be^{-3t}$ must be function of t, not x</p> <p>P.I. $x = pt + q$ (or $x = at^2 + bt + c$)</p> <p>$4p + 3(pt + q) = kt + 5$ $3p = k$ (Form at least one eqn. in p and/or q)</p> <p style="padding-left: 100px;">$4p + 3q = 5$</p> <p>$p = \frac{k}{3},$ $q = \frac{5}{3} - \frac{4k}{9}$ $\left(= \frac{15 - 4k}{9} \right)$</p> <p>General solution: $x = Ae^{-t} + Be^{-3t} + \frac{kt}{3} + \frac{15 - 4k}{9}$ must include $x =$ and be function of t</p> <p>(b) When $k = 6,$ $x = 2t - 1$</p>	<p>M1 A1</p> <p>A1</p> <p>B1</p> <p>M1</p> <p>A1</p> <p>A1 ft (7)</p> <p>M1 A1cao (2)</p> <p style="text-align: right;">9</p>
	<p>(a) M1 for auxiliary equation substantially correct B1 not awarded for $x = kt + \text{constant}$</p> <p>(b) M mark for using $k = 6$ to derive a linear expression in t. (cf must have involved negative exponentials only) so e.g. $y = 2t - 1$ is M1 A0</p>	

Question number	Scheme	Marks
5.	<p>(a) $\frac{4}{x} = \frac{x}{2} + 3$ $x^2 + 6x - 8 = 0$ $x = \dots, \left(\frac{-6 \pm \sqrt{68}}{2} \right)$ $-3 \pm \sqrt{17}$ - root not needed</p> <p>$-\frac{4}{x} = \frac{x}{2} + 3,$ $x^2 + 6x + 8 = 0$ $x = -4$ and -2</p> <p>Three correct solutions (and no extras): $-4, -2, -3 + \sqrt{17}$</p> <p>(b)  Line through point on -ve x axis and + y axis Curve 3 Intersections in correct quadrants</p> <p>(c) $-4 < x < -2,$ $x > -3 + \sqrt{17}$ o.e.</p>	<p>M1, A1</p> <p>M1, A1</p> <p>A1 (5)</p> <p>B1 B1 B1 (3)</p> <p>B1, B1 (2)</p> <p>10</p>
	<p>(a) <u>Alternative using squaring method</u> Square both sides and attempt to find roots $x^4 + 12x^3 + 36x^2 - 64 = 0$ gives $x = -2$ and $x = -4$ Obtain quadratic factor, divide find solutions of quadratic and obtain $(-3 \pm \sqrt{17})$ Last mark as before</p> <p>(c) Use of \leq instead of $<$ lose last B1 Extra inequalities lose last B1</p>	<p>M1 A1 M1 A1</p>

Question number	Scheme	Marks
6.	<p>(a) $\frac{2}{(r+1)(r+3)} = \frac{1}{r+1} - \frac{1}{r+3}$ M: $\frac{2}{(r+1)(r+3)} = \frac{A}{r+1} + \frac{B}{r+3}$</p> <p>(b) $r = 1: \left(\frac{2}{2 \times 4}\right) = \frac{1}{2} - \frac{1}{4}$</p> <p>$r = 2: \left(\frac{2}{3 \times 5}\right) = \frac{1}{3} - \frac{1}{5}$</p> <p>... $r = n - 1: \left(\frac{2}{n(n+2)}\right) = \frac{1}{n} - \frac{1}{n+2}$</p> <p>$r = n: \left(\frac{2}{(n+1)(n+3)}\right) = \frac{1}{n+1} - \frac{1}{n+3}$</p> <p>Summing: $\sum = \frac{1}{2} + \frac{1}{3} - \frac{1}{n+2} - \frac{1}{n+3}$</p> <p>$= \frac{5(n+2)(n+3) - 6(n+3) - 6(n+2)}{6(n+2)(n+3)} = \frac{n(5n+13)}{6(n+2)(n+3)}$</p> <p>(c) $\sum_{21}^{30} = \sum_1^{30} - \sum_1^{20} = \frac{30 \times 163}{6 \times 32 \times 33} - \frac{20 \times 113}{6 \times 22 \times 23}, \quad = 0.02738$</p>	<p>M1 A1 (2)</p> <p>M1</p> <p>A1 ft</p> <p>M1 A1</p> <p>d M1 A1cso (6)</p> <p>M1A1ft,A1cso (3)</p> <p>(11)</p>
	<p>(b) The first M1 requires list of first two and last two terms The A1 must be correct but ft on their A and B The second M1 requires terms to be eliminated and A1 is cao</p> <p>(c) The M mark is also allowed for $\sum_1^{30} - \sum_1^{21}$ applied with numbers included</p> <p>Using $u_{30} - u_{20}$ scores M0 A0 A0</p> <p>The first A1 is ft their A and B or could include A and B, but final A1 is cao but accept 0.027379775599 to 5 or more decimal places..</p>	

Question number	Scheme	Marks
7.	<p>(a) $\frac{dy}{dx} = v + x \frac{dv}{dx}$</p> $\left(v + x \frac{dv}{dx}\right) = \frac{x}{vx} + \frac{3vx}{x} \Rightarrow x \frac{dv}{dx} = 2v + \frac{1}{v} \quad (*)$ <p>(b) $\int \frac{v}{1+2v^2} dv = \int \frac{1}{x} dx$</p> $\frac{1}{4} \ln(1+2v^2), = \ln x (+C)$ $Ax^4 = 1 + 2v^2$ $Ax^4 = 1 + 2\left(\frac{y}{x}\right)^2 \text{ so } y = \sqrt{\frac{Ax^6 - x^2}{2}} \text{ or } y = x\sqrt{\frac{Ax^4 - 1}{2}} \text{ or } y = x\sqrt{\left(\frac{1}{2}e^{4\ln x + 4c} - \frac{1}{2}\right)}$ <p>(c) $x = 1$ at $y = 3$: $3 = \sqrt{\frac{A-1}{2}} \quad A = \dots$</p> $y = \sqrt{\frac{19x^6 - x^2}{2}} \text{ or } y = x\sqrt{\frac{19x^4 - 1}{2}}$	<p>B1</p> <p>M1 A1 (3)</p> <p>M1</p> <p>dM1 A1, B1</p> <p>d M1</p> <p>M1 A1 (7)</p> <p>M1</p> <p>A1 (2) 12</p>
	<p>(a) B1 for statement printed or for $\frac{dy}{dx} = \left(x + v \frac{dx}{dv}\right) \frac{dv}{dx}$</p> <p>First M1 is for RHS of equation only but for A1 need whole answer correct .</p> <p>(b) First M1 accept $\int \frac{1}{2v + \frac{1}{v}} dv = \int \frac{1}{x} dx$</p> <p>Second M1 requires an integration of correct form $\frac{1}{4}$ may be missing</p> <p>A1 for LHS correct with $\frac{1}{4}$ and B1 is independent and is for $\ln x$</p> <p>Third M1 is dependent and needs correct application of log laws</p> <p>Fourth M1 is independent and merely requires return to y/x for v</p> <p>N.B. There is an IF method possible after suitable rearrangement – see note.</p>	

Question number	Scheme	Marks
8.	<p>(a) $r \cos \theta = 4(\cos \theta - \cos^2 \theta)$ or $r \cos \theta = 4 \cos \theta - 2 \cos 2\theta - 2$</p> $\frac{d(r \cos \theta)}{d\theta} = 4(-\sin \theta + 2 \cos \theta \sin \theta) \text{ or } \frac{d(r \cos \theta)}{d\theta} = 4(-\sin \theta + \sin 2\theta)$ $4(-\sin \theta + 2 \cos \theta \sin \theta) = 0 \Rightarrow \cos \theta = \frac{1}{2} \text{ which is satisfied by } \theta = \frac{\pi}{3} \text{ and } r = 2(*)$ <p>(b) $\frac{1}{2} \int r^2 d\theta = (8) \int (1 - 2 \cos \theta + \cos^2 \theta) d\theta$</p> $= (8) \left[\theta - 2 \sin \theta + \frac{\sin 2\theta}{4} + \frac{\theta}{2} \right]$ $8 \left[\frac{3\theta}{2} - 2 \sin \theta + \frac{\sin 2\theta}{4} \right]_{-\pi/3}^{\pi/2} = 8 \left(\left(\frac{3\pi}{4} - 2 \right) - \left(\frac{\pi}{2} - \sqrt{3} + \frac{\sqrt{3}}{8} \right) \right) = 2\pi - 16 + 7\sqrt{3}$ <p>Triangle: $\frac{1}{2}(r \cos \theta)(r \sin \theta) = \frac{1}{2} \times 1 \times \sqrt{3} = \frac{\sqrt{3}}{2}$</p> <p>Total area: $(2\pi - 16 + 7\sqrt{3}) + \frac{\sqrt{3}}{2} = (2\pi - 16) + \frac{15\sqrt{3}}{2}$</p>	<p>B1</p> <p>M1 A1</p> <p>d M1 A1 (5)</p> <p>M1</p> <p>M1 A1</p> <p>M1</p> <p>M1 A1</p> <p>(A1) A1 (8)</p> <p>13</p>
	<p>(a) <u>Alternative for first 3 marks:</u></p> $\frac{dr}{d\theta} = 4 \sin \theta \quad \text{B1}$ $\frac{dx}{d\theta} = -r \sin \theta + \cos \theta \frac{dr}{d\theta} = -4 \sin \theta + 8 \sin \theta \cos \theta \quad \text{M1 A1}$ <p>Substituting $r = 2$ and $\theta = \frac{\pi}{3}$ into original equation scores 0 marks.</p> <p>(b) M1 needs attempt to expand $(1 - \cos \theta)^2$ giving three terms (allow slips)</p> <p>Second M1 needs integration of $\cos^2 \theta$ using $\cos 2\theta \pm 1$</p> <p>Third M1 needs correct limits- may evaluate two areas and subtract</p> <p>M1 needs attempt at area of triangle and A1 for cao</p> <p>Next A1 is for value of area within curve, then final A1 is cao, must be exact but allow 4 terms and isw for incorrect collection of terms</p> <p>Special case for use of $r \sin \theta$ gives B0M1A0M0A0</p>	