

1. A random sample X_1, X_2, \dots, X_{10} is taken from a population with mean μ and variance σ^2 .

(a) Determine the bias, if any, of each of the following estimators of μ .

$$\theta_1 = \frac{X_3 + X_4 + X_5}{3},$$

$$\theta_2 = \frac{X_{10} - X_1}{3},$$

$$\theta_3 = \frac{3X_1 + 2X_2 + X_{10}}{6}.$$

(4)

(b) Find the variance of each of these estimators.

(5)

(c) State, giving reasons, which of these three estimators for μ is

(i) the best estimator,

(ii) the worst estimator.

(4)



2. A large number of students are split into two groups A and B . The students sit the same test but under different conditions. Group A has music playing in the room during the test, and group B has no music playing during the test. Small samples are then taken from each group and their marks recorded. The marks are normally distributed.

The marks are as follows:

Sample from Group A	42	40	35	37	34	43	42	44	49
Sample from Group B	40	44	38	47	38	37	33		

- (a) Stating your hypotheses clearly, and using a 10% level of significance, test whether or not there is evidence of a difference between the variances of the marks of the two groups. **(8)**

- (b) State clearly an assumption you have made to enable you to carry out the test in part (a). **(1)**

- (c) Use a two tailed test, with a 5% level of significance, to determine if the playing of music during the test has made any difference in the mean marks of the two groups. State your hypotheses clearly. **(7)**

- (d) Write down what you can conclude about the effect of music on a student's performance during the test. **(1)**





Leave
blank

Question 2 continued

Lined writing area for the answer to Question 2.



Turn over



H 3 1 3 2 7 A 0 7 2 4



Question 2 continued

Leave
blank

Lined area for writing the answer to Question 2.





Question 5 continued

Leave
blank

Lined writing area for the answer to Question 5.

(Total 8 marks)

Q5



6. A drug is claimed to produce a cure to a certain disease in 35% of people who have the disease. To test this claim a sample of 20 people having this disease is chosen at random and given the drug. If the number of people cured is between 4 and 10 inclusive the claim will be accepted. Otherwise the claim will not be accepted.

(a) Write down suitable hypotheses to carry out this test. (2)

(b) Find the probability of making a Type I error. (3)

The table below gives the value of the probability of the Type II error, to 4 decimal places, for different values of p where p is the probability of the drug curing a person with the disease.

P(cure)	0.2	0.3	0.4	0.5
P(Type II error)	0.5880	r	0.8565	s

(c) Calculate the value of r and the value of s . (3)

(d) Calculate the power of the test for $p = 0.2$ and $p = 0.4$ (2)

(e) Comment, giving your reasons, on the suitability of this test procedure. (2)



7. An engineering firm buys steel rods. The steel rods from its present supplier are known to have a mean tensile strength of 230 N/mm².

A new supplier of steel rods offers to supply rods at a cheaper price than the present supplier. A random sample of ten rods from this new supplier gave tensile strengths, x N/mm², which are summarised below.

Sample size	Σx	Σx^2
10	2283	524079

- (a) Stating your hypotheses clearly, and using a 5% level of significance, test whether or not the rods from the new supplier have a tensile strength lower than the present supplier. (You may assume that the tensile strength is normally distributed). **(7)**

- (b) In the light of your conclusion to part (a) write down what you would recommend the engineering firm to do. **(1)**



BLANK PAGE

