

Mark Scheme (Results)

November 2012

GCSE Chemistry  
5CH2H/01

## **Edexcel and BTEC Qualifications**

Edexcel and BTEC qualifications come from Pearson, the world's leading learning company. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at [www.edexcel.com](http://www.edexcel.com) or [www.btec.co.uk](http://www.btec.co.uk) for our BTEC qualifications.

Alternatively, you can get in touch with us using the details on our contact us page at [www.edexcel.com/contactus](http://www.edexcel.com/contactus).

If you have any subject specific questions about this specification that require the help of a subject specialist, you can speak directly to the subject team at Pearson.

Their contact details can be found on this link: [www.edexcel.com/teachingservices](http://www.edexcel.com/teachingservices).

You can also use our online Ask the Expert service at [www.edexcel.com/ask](http://www.edexcel.com/ask). You will need an Edexcel username and password to access this service.

### **Pearson: helping people progress, everywhere**

Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: [www.pearson.com/uk](http://www.pearson.com/uk)

November 2012

Publications Code UG034057

All the material in this publication is copyright

© Pearson Education Ltd 2012


GCSE Chemistry 5CH2H/01 Mark Scheme – November 2012

Question Number	Answer	Acceptable answers	Mark
<b>1(a)</b>	C : copper sulfate and sodium chloride		<b>(1)</b>

Question Number	Answer	Acceptable answers	Mark
<b>1(b)</b>	copper sulfate (1) blue-green (1) or sodium chloride (1) yellow (1) colour mark consequential on correct metal (compound)	allow blue or green or green-blue  reject orange and yellow-orange	<b>(2)</b>

Question Number	Answer	Acceptable answers	Mark
<b>1(c)(i)</b>	An explanation linking  weak <b>intermolecular</b> forces /weak forces between <b>molecules</b> (1)  little {heat / energy} needed to separate (molecules) (1)	bonds / attractions in place of forces  intermolecular forces between {atoms / bonds} loses 1 <sup>st</sup> marking point  any answer in terms of covalent or ionic bonding scores zero	<b>(2)</b>

Question Number	Answer	Acceptable answers	Mark
<b>1(c)(ii)</b>	A description linking  use separating funnel (1)  run off lower {layer / liquid} / OWTTE (1)	alternative description of separating funnel eg funnel with a tap at the bottom suitable labelled diagram burette  allow layers / liquids to separate  ignore fractional distillation	<b>(2)</b>

Question Number	Answer	Acceptable answers	Mark
1(d)	 <p data-bbox="336 533 807 640">shared pair in molecule (1) rest of molecule consequent on first mark (1)</p>	<p data-bbox="868 349 1331 421">Allow a diagram without labels for 2 marks</p> <p data-bbox="868 600 1299 669">any symbols shown must be correct for the 2<sup>nd</sup> mark</p> <p data-bbox="868 707 1321 777">allow any combination of dots and crosses for electrons</p> <p data-bbox="868 815 1342 844">wrong compound = zero marks</p>	(2)

Question Number	Answer	Acceptable answers	Mark
<b>2(a)(i)</b>	soft / low melting point / low boiling point	<b>easily</b> cut with a knife = soft low density malleable solid at room temp.  ignore float on water reject chemical properties	<b>(1)</b>

Question Number	Answer	Acceptable answers	Mark
<b>2(a)(ii)</b>	An explanation linking  (all have) one electron in outer shell (2)	one outer electron = 2 marks  group number shows number of electrons in outer shell = 2 marks  same number of electrons in outer shell = 1 mark  incorrect number of electrons in the outer shell = 1 mark  accept outer orbit / highest energy level in place of outer shell	<b>(2)</b>

Question Number	Answer	Acceptable answers	Mark
<b>2(b)(i)</b>	A description including any two of effervescence / fizzing / bubbles (1) potassium floats (1) moves (on surface) (1) potassium forms ball / melts (1) potassium decreases in size / disappears / dissolves (1) (lilac) flame / catches fire (1) spits / explodes / sparks (1)	ignore ignites ignore smoke	<b>(2)</b>

Question Number	Answer	Acceptable answers	Mark
<b>2(b)(ii)</b>	D : $2K + 2H_2O \rightarrow 2KOH + H_2$		<b>(1)</b>

Question Number	Answer	Acceptable answers	Mark
<b>2(c)</b>	An explanation linking any two of increasing {size /radius (of atom) / number of shells} (1) increased shielding (of outer electron) (1) less attraction for (outer) electron (1)	easier to remove (outer) electron	<b>(2)</b>

Question Number	Answer	Acceptable answers	Mark
<b>3(a)(i)</b>	A, B and C	Mg Ca Au (any order) magnesium calcium gold (any order)	<b>(1)</b>

Question Number	Answer	Acceptable answers	Mark
<b>3(a)(ii)</b>	A and B	Mg Ca (any order) magnesium calcium (any order)	<b>(1)</b>

Question Number	Answer	Acceptable answers	Mark
<b>3(b)</b>	8 (protons)		<b>(1)</b>

Question Number	Answer	Acceptable answers	Mark
<b>3(c)(i)</b>	A : 10		<b>(1)</b>

Question Number	Answer	Acceptable answers	Mark
<b>3(c)(ii)</b>	(in 100 atoms)  mass of mass number 20 atoms = $20 \times 90$ (1) mass of mass number 22 atoms = $22 \times 10$ (1) relative atomic mass = $\{(22 \times 10) + (20 \times 90)\} / 100$ (=20.2) (1)  OR  20 contributes = $90/100 \times 20$ (1) 22 contributes = $10/100 \times 22$ (1) relative atomic mass = $90/100 \times 20 + 10/100 \times 22$ (= 20.2) (1)	20.2 = 3 marks  21.8 = 2 marks (only 1 error made)	<b>(3)</b>

Question Number	Answer	Acceptable answers	Mark
<b>3(d)</b>	An explanation linking any two of  (the element is) group 0 / noble gas / unreactive / inert / does not react (1) { (has) 8 electrons / full } outer shell (1) prevents filament from reacting (1)	ignore 'not very reactive'  does not {gain / lose / share} electrons	<b>(2)</b>



Question Number	Answer	Acceptable answers	Mark
<b>4(a)</b>	to allow air/oxygen in	to ensure magnesium reacts/burns / combusts	<b>(1)</b>

Question Number	Answer	Acceptable answers	Mark
<b>4(b)(i)</b>	all points correctly plotted to half a small square (2) line of best fit (1)	Allow one mark for four or five correctly plotted points ecf their points	<b>(3)</b>

Question Number	Answer	Acceptable answers	Mark
<b>4(b)(ii)</b>	Any one from  not all magnesium {burned / reacted} / some left / incomplete reaction not enough air/oxygen some magnesium oxide / smoke lost	lid not lifted / not enough times lid left off too long (so loses MgO)	<b>(1)</b>

Question Number	Answer	Acceptable answers	Mark
<b>4(c)</b>	$2\text{Mg} + \text{O}_2 \rightarrow 2\text{MgO}$ left hand formulae (1) right hand formula (1) balancing correct formulae (1)	correct multiples	<b>(3)</b>

Question Number	Answer	Acceptable answers	Mark
<b>4(d)</b>	0.414 / 207 or 0.064 / 16 (1) 0.002 : 0.004 or 1 : 2 (1) empirical formula $\text{PbO}_2$ (1)	if 207 / 0.414 and 16 / 0.064 ratio 500 : 250 or 2 : 1 (1) empirical formula $\text{Pb}_2\text{O}$ (1)  allow 3 marks for 0.414 / 207 or 0.064 / 32 ratio 1 : 1 empirical formula $\text{PbO}_2$  allow 2 marks for if 0.414 / 207 and 0.064 / 32 ratio 1 : 1 empirical formula $\text{PbO}$	<b>(3)</b>

Question Number	Answer	Acceptable answers	Mark
<b>5(a)</b>	An explanation linking two of the following  temperature decreases (1) {heat / energy} taken in (1) (so process) endothermic (1)	ignore references to bond breaking / making  heat given out / exothermic = 1 max.	<b>(2)</b>

Question Number	Answer	Acceptable answers	Mark
<b>5(b)</b>	Shown correctly on diagram:  horizontal line to right of reactant (1) product line below reactant line (1)	ignore any connecting lines product label not needed	<b>(2)</b>

Question Number	Answer	Acceptable answers	Mark
<b>5(c)</b>	D : heat energy is required heat energy is released		<b>(1)</b>

Question Number		Indicative Content	Mark
<b>QWC</b>	<b>*5(d)</b>	<p>An explanation including some of the following points</p> <p><b>smaller pieces of solid</b>  of same mass  larger surface area  more frequent collisions  higher rate of reaction</p> <p><b>higher temperature</b>  particles move faster  more frequent collisions  particles have more energy  more collisions have required energy to react / activation energy  more collisions successful  higher rate of reaction</p> <p>ORA</p>	<b>(6)</b>
<b>Level I</b>	<b>0</b>	No rewardable content	
<b>1</b>	<b>1 - 2</b>	<p>a limited explanation of one of factors e.g. at higher temperature higher rate  e.g. when particles smaller size higher rate  the answer communicates ideas using simple language and uses limited scientific terminology  spelling, punctuation and grammar are used with limited accuracy</p>	
<b>2</b>	<b>3 - 4</b>	<p>a simple explanation e.g. at higher temperature particles move faster, more collisions so higher rate  e.g. smaller sized particles (of same mass) have greater surface area so higher rate  the answer communicates ideas showing some evidence of clarity and organisation and uses scientific terminology appropriately  spelling, punctuation and grammar are used with some accuracy</p>	
<b>3</b>	<b>5 - 6</b>	<p>a detailed explanation e.g. (when particles collide they) only react when they have sufficient energy/activation energy and at a higher temperature more of the particles have sufficient energy/activation energy so more collisions will be successful and when particles smaller size higher rate  the answer communicates ideas clearly and coherently uses a range of scientific terminology accurately  spelling, punctuation and grammar are used with few errors</p>	

Question Number	Answer	Acceptable answers	Mark
<b>6(a)</b>	D : $\text{Ca}(\text{NO}_3)_2$		<b>(1)</b>

Question Number	Answer	Acceptable answers	Mark
<b>6(b)</b>	C : 8		<b>(1)</b>

Question Number	Answer	Acceptable answers	Mark
<b>6(c)</b>	<p>Description including four of the following</p> <p>sodium - 2.8.1 / 1 electron in outer shell (1)  sodium (atoms) lose electrons (1)  one per atom (1)  (forms) <math>\text{Na}^+</math> (1)  sulphur - 2.8.6 / 6 electrons in outer shell (1)  sulfur (atoms) gain electrons (1)  two per atom (1)  (forms) <math>\text{S}^{2-}</math> (1)  two sodium atoms / ions combine with one sulfur atom / ion (1)  formula is <math>\text{Na}_2\text{S}</math> (1)</p>	<p>Marks can be gained using diagrams</p> <p>mention of shared electrons / covalent bonding in words or diagram = max 2 marks</p>	<b>(4)</b>

Question Number		Indicative Content	Mark
<b>QWC</b>	<b>* 6(d)</b>	<p>A description including some of the following points</p> <p><b>solid</b>  {regular arrangement/ lattice} (of ions)  sodium/Na<sup>+</sup> ions  chloride /Cl<sup>-</sup> ions  (held together by)  strong (ionic) bonds  strong (electrostatic) forces of attraction  between oppositely charged ions / positive and negatively charged ions  closely packed together  (when solid) does not conduct  because ions cannot move</p> <p><b>molten</b>  heat energy {overcomes/breaks} (strong ionic) bonds  strong (electrostatic) forces of attraction  between oppositely charged ions / positive and negatively charged ions  ions can move  (therefore) conducts when molten</p>	<b>(6)</b>
<b>Level</b>	<b>0</b>	No rewardable content	
<b>1</b>	<b>1 - 2</b>	a limited explanation e.g. does not conduct when solid e.g. does conduct when molten the answer communicates ideas using simple language and uses limited scientific terminology spelling, punctuation and grammar are used with limited accuracy	
<b>2</b>	<b>3 - 4</b>	a simple explanation e.g. does not conduct when solid, does conduct when molten because {ions / particles / atoms} can move the answer communicates ideas showing some evidence of clarity and organisation and uses scientific terminology appropriately spelling, punctuation and grammar are used with some accuracy	
<b>3</b>	<b>5 - 6</b>	a detailed explanation e.g. solid has strong ionic bonds (between oppositely charged ions), does not conduct when solid because ions cannot move, does conduct when molten because ions can move the answer communicates ideas clearly and coherently uses a range of scientific terminology accurately spelling, punctuation and grammar are used with few errors	

Further copies of this publication are available from  
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467

Fax 01623 450481

Email [publication.orders@edexcel.com](mailto:publication.orders@edexcel.com)

Order Code UG034057 November 2012

For more information on Edexcel qualifications, please visit our website  
[www.edexcel.com](http://www.edexcel.com)

Pearson Education Limited. Registered company number 872828  
with its registered office at Edinburgh Gate, Harlow, Essex CM20 2JE

Ofqual  




Llywodraeth Cymru  
Welsh Assembly Government



Rewarding Learning