| Centre No. | | | | | | Paper Reference | | | | | | | Surname | Initial(s) |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| Candidate No. | | | | | | 6 | 6 | 6 | 6 | / | 0 | 1 | Signature | |

Paper Reference(s)

**6666/01**

# Edexcel GCE

## Core Mathematics C4

## Advanced

Monday 28 January 2013 – Morning

Time: 1 hour 30 minutes

Examiner's use only

| | | |
|---|---|---|
| | | |

Team Leader's use only

| | | |
|---|---|---|
| | | |

| Question Number | Leave Blank |
|---|---|
| 1 | |
| 2 | |
| 3 | |
| 4 | |
| 5 | |
| 6 | |
| 7 | |
| 8 | |
| Total | |

**Materials required for examination**
Mathematical Formulae (Pink)

**Items included with question papers**
Nil

**Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation or symbolic differentiation/integration, or have retrievable mathematical formulae stored in them.**

### Instructions to Candidates

In the boxes above, write your centre number, candidate number, your surname, initials and signature.
Check that you have the correct question paper.
Answer ALL the questions.
You must write your answer for each question in the space following the question.
When a calculator is used, the answer should be given to an appropriate degree of accuracy.

### Information for Candidates

A booklet 'Mathematical Formulae and Statistical Tables' is provided.
Full marks may be obtained for answers to ALL questions.
The marks for individual questions and the parts of questions are shown in round brackets: e.g. **(2)**.
There are 8 questions in this question paper. The total mark for this paper is 75.
There are 28 pages in this question paper. Any blank pages are indicated.

### Advice to Candidates

You must ensure that your answers to parts of questions are clearly labelled.
You should show sufficient working to make your methods clear to the Examiner.
Answers without working may not gain full credit.

This publication may be reproduced only in accordance with Pearson Education Ltd copyright policy.
©2013 Pearson Education Ltd.

Printer's Log. No.
P41860A
W850/R6666/57570 5/5/5/6/

P 4 1 8 6 0 A 0 1 2 8

*Turn over*

PEARSON

Leave blank

**1.** Given

$$f(x) = (2 + 3x)^{-3}, \qquad |x| < \frac{2}{3}$$

find the binomial expansion of f($x$), in ascending powers of $x$, up to and including the term in $x^3$.

Give each coefficient as a simplified fraction.

**(5)**



P 4 1 8 6 0 A 0 2 2 8

Leave blank

**Question 1 continued**

**Q1**

**(Total 5 marks)**



P 4 1 8 6 0 A 0 3 2 8

**Turn over**

Leave blank

**2.** (a) Use integration to find

$$\int \frac{1}{x^3} \ln x \, \mathrm{d}x$$

**(5)**

(b) Hence calculate

$$\int_1^2 \frac{1}{x^3} \ln x \, \mathrm{d}x$$

**(2)**



P 4 1 8 6 0 A 0 4 2 8

Leave blank

**Question 2 continued**

**Q2**

**(Total 7 marks)**

Leave blank

**3.** Express $\dfrac{9x^2 + 20x - 10}{(x + 2)(3x - 1)}$ in partial fractions.

**(4)**

Leave blank

**Question 3 continued**

**Q3**

**(Total 4 marks)**

Leave blank

**4.**




**Figure 1**

Figure 1 shows a sketch of part of the curve with equation $y = \dfrac{x}{1 + \sqrt{x}}$. The finite region $R$, shown shaded in Figure 1, is bounded by the curve, the $x$-axis, the line with equation $x = 1$ and the line with equation $x = 4$.

(a) Complete the table with the value of $y$ corresponding to $x = 3$, giving your answer to 4 decimal places.

**(1)**

| $x$ | 1 | 2 | 3 | 4 |
|---|---|---|---|---|
| $y$ | 0.5 | 0.8284 | | 1.3333 |

(b) Use the trapezium rule, with all the values of $y$ in the completed table, to obtain an estimate of the area of the region $R$, giving your answer to 3 decimal places.

**(3)**

(c) Use the substitution $u = 1 + \sqrt{x}$, to find, by integrating, the exact area of $R$.

**(8)**



P 4 1 8 6 0 A 0 8 2 8

Leave blank

**Question 4 continued**

Leave blank

**Question 4 continued**

Leave blank

**Question 4 continued**

**Q4**

**(Total 12 marks)**

Leave blank

**5.**




**Figure 2**

Figure 2 shows a sketch of part of the curve $C$ with parametric equations

$$x = 1 - \frac{1}{2}t, \quad y = 2^t - 1$$

The curve crosses the $y$-axis at the point $A$ and crosses the $x$-axis at the point $B$.

(a) Show that $A$ has coordinates (0, 3).

**(2)**

(b) Find the $x$ coordinate of the point $B$.

**(2)**

(c) Find an equation of the normal to $C$ at the point $A$.

**(5)**

The region $R$, as shown shaded in Figure 2, is bounded by the curve $C$, the line $x = -1$ and the $x$-axis.

(d) Use integration to find the exact area of $R$.

**(6)**



P 4 1 8 6 0 A 0 1 2 2 8

Leave blank

**Question 5 continued**

Leave blank

**Question 5 continued**

Leave blank

**Question 5 continued**

**Q5**

**(Total 15 marks)**



P 4 1 8 6 0 A 0 1 5 2 8

**Turn over**

Leave blank

**6.**




**Figure 3**

Figure 3 shows a sketch of part of the curve with equation $y = 1 - 2\cos x$, where $x$ is measured in radians. The curve crosses the $x$-axis at the point $A$ and at the point $B$.

(a) Find, in terms of $\pi$, the $x$ coordinate of the point $A$ and the $x$ coordinate of the point $B$.
**(3)**

The finite region $S$ enclosed by the curve and the $x$-axis is shown shaded in Figure 3. The region $S$ is rotated through $2\pi$ radians about the $x$-axis.

(b) Find, by integration, the exact value of the volume of the solid generated.
**(6)**



P 4 1 8 6 0 A 0 1 6 2 8

Leave blank

**Question 6 continued**

Leave blank

**Question 6 continued**

Leave blank

**Question 6 continued**

**Q6**

**(Total 9 marks)**



P 4 1 8 6 0 A 0 1 9 2 8

**Turn over**

Leave blank

**7.** With respect to a fixed origin $O$, the lines $l_1$ and $l_2$ are given by the equations

$$l_1 : \mathbf{r} = (9\mathbf{i} + 13\mathbf{j} - 3\mathbf{k}) + \lambda(\mathbf{i} + 4\mathbf{j} - 2\mathbf{k})$$

$$l_2 : \mathbf{r} = (2\mathbf{i} - \mathbf{j} + \mathbf{k}) + \mu(2\mathbf{i} + \mathbf{j} + \mathbf{k})$$

where $\lambda$ and $\mu$ are scalar parameters.

(a) Given that $l_1$ and $l_2$ meet, find the position vector of their point of intersection.

**(5)**

(b) Find the acute angle between $l_1$ and $l_2$, giving your answer in degrees to 1 decimal place.

**(3)**

Given that the point $A$ has position vector $4\mathbf{i} + 16\mathbf{j} - 3\mathbf{k}$ and that the point $P$ lies on $l_1$ such that $AP$ is perpendicular to $l_1$,

(c) find the exact coordinates of $P$.

**(6)**



P 4 1 8 6 0 A 0 2 0 2 8

Leave blank

**Question 7 continued**

Leave blank

**Question 7 continued**

Leave blank

**Question 7 continued**

**(Total 14 marks)**

**Q7**

Leave blank

**8.** A bottle of water is put into a refrigerator. The temperature inside the refrigerator remains constant at 3 °C and $t$ minutes after the bottle is placed in the refrigerator the temperature of the water in the bottle is $\theta$ °C.

The rate of change of the temperature of the water in the bottle is modelled by the differential equation,

$$\frac{d\theta}{dt} = \frac{(3 - \theta)}{125}$$

(a) By solving the differential equation, show that,

$$\theta = Ae^{-0.008t} + 3$$

where $A$ is a constant.

**(4)**

Given that the temperature of the water in the bottle when it was put in the refrigerator was 16 °C,

(b) find the time taken for the temperature of the water in the bottle to fall to 10 °C, giving your answer to the nearest minute.

**(5)**

Leave blank

**Question 8 continued**

Leave blank

**Question 8 continued**

Leave blank

**Question 8 continued**

Leave blank

**Question 8 continued**

Q8

**(Total 9 marks)**

**TOTAL FOR PAPER: 75 MARKS**

**END**



P 4 1 8 6 0 A 0 2 8 2 8