








2.



**Figure 1**

A non-uniform rod  $AB$  has length 3 m and mass 4.5 kg. The rod rests in equilibrium, in a horizontal position, on two smooth supports at  $P$  and at  $Q$ , where  $AP = 0.8$  m and  $QB = 0.6$  m, as shown in Figure 1. The centre of mass of the rod is at  $G$ . Given that the magnitude of the reaction of the support at  $P$  on the rod is twice the magnitude of the reaction of the support at  $Q$  on the rod, find

(a) the magnitude of the reaction of the support at  $Q$  on the rod, **(3)**

(b) the distance  $AG$ . **(4)**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



**Question 2 continued**

Lined area for writing the answer to Question 2.

**(Total 7 marks)**

**Q2**



P 4 0 6 8 9 A 0 5 2 8

3.




Figure 2

A box of mass 5 kg lies on a rough plane inclined at  $30^\circ$  to the horizontal. The box is held in equilibrium by a horizontal force of magnitude 20 N, as shown in Figure 2. The force acts in a vertical plane containing a line of greatest slope of the inclined plane. The box is in equilibrium and on the point of moving down the plane. The box is modelled as a particle.

Find

- (a) the magnitude of the normal reaction of the plane on the box, (4)

- (b) the coefficient of friction between the box and the plane. (5)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---











4. A car is moving on a straight horizontal road. At time  $t = 0$ , the car is moving with speed  $20 \text{ m s}^{-1}$  and is at the point  $A$ . The car maintains the speed of  $20 \text{ m s}^{-1}$  for 25 s. The car then moves with constant deceleration  $0.4 \text{ m s}^{-2}$ , reducing its speed from  $20 \text{ m s}^{-1}$  to  $8 \text{ m s}^{-1}$ . The car then moves with constant speed  $8 \text{ m s}^{-1}$  for 60 s. The car then moves with constant acceleration until it is moving with speed  $20 \text{ m s}^{-1}$  at the point  $B$ .

(a) Sketch a speed-time graph to represent the motion of the car from  $A$  to  $B$ . (3)

(b) Find the time for which the car is decelerating. (2)

Given that the distance from  $A$  to  $B$  is 1960 m,

(c) find the time taken for the car to move from  $A$  to  $B$ . (8)





**Question 4 continued**

A series of horizontal lines provided for writing an answer to Question 4.



### Question 4 continued

Ruled area for writing the answer to Question 4.

**(Total 13 marks)**

**Q4**











Leave  
blank

### Question 5 continued

Handwritten response area consisting of 30 horizontal lines.

(Total 12 marks)

Q5



6. [In this question  $\mathbf{i}$  and  $\mathbf{j}$  are horizontal unit vectors due east and due north respectively and position vectors are given with respect to a fixed origin.]

A ship  $S$  is moving with constant velocity  $(-12\mathbf{i} + 7.5\mathbf{j}) \text{ km h}^{-1}$ .

- (a) Find the direction in which  $S$  is moving, giving your answer as a bearing. (3)

At time  $t$  hours after noon, the position vector of  $S$  is  $\mathbf{s}$  km. When  $t = 0$ ,  $\mathbf{s} = 40\mathbf{i} - 6\mathbf{j}$ .

- (b) Write down  $\mathbf{s}$  in terms of  $t$ . (2)

A fixed beacon  $B$  is at the point with position vector  $(7\mathbf{i} + 12.5\mathbf{j}) \text{ km}$ .

- (c) Find the distance of  $S$  from  $B$  when  $t = 3$  (4)

- (d) Find the distance of  $S$  from  $B$  when  $S$  is due north of  $B$ . (4)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---









7.



**Figure 3**

Two particles  $P$  and  $Q$ , of mass 0.3 kg and 0.5 kg respectively, are joined by a light horizontal rod. The system of the particles and the rod is at rest on a horizontal plane. At time  $t = 0$ , a constant force  $\mathbf{F}$  of magnitude 4 N is applied to  $Q$  in the direction  $PQ$ , as shown in Figure 3. The system moves under the action of this force until  $t = 6$  s. During the motion, the resistance to the motion of  $P$  has constant magnitude 1 N and the resistance to the motion of  $Q$  has constant magnitude 2 N.

Find

- (a) the acceleration of the particles as the system moves under the action of  $\mathbf{F}$ , (3)
- (b) the speed of the particles at  $t = 6$  s, (2)
- (c) the tension in the rod as the system moves under the action of  $\mathbf{F}$ . (3)

At  $t = 6$  s,  $\mathbf{F}$  is removed and the system decelerates to rest. The resistances to motion are unchanged. Find

- (d) the distance moved by  $P$  as the system decelerates, (4)
- (e) the thrust in the rod as the system decelerates. (3)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---









**Question 7 continued**

Lined area for writing the answer to Question 7.







**Question 7 continued**

Handwriting practice area consisting of multiple horizontal lines for text entry.

**Q7**

--	--

**(Total 15 marks)**

**TOTAL FOR PAPER: 75 MARKS**

**END**

