

# Mark Scheme (Results) January 2010

GCE

## Mechanics M2 (6678)

Edexcel is one of the leading examining and awarding bodies in the UK and throughout the world. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. Through a network of UK and overseas offices, Edexcel's centres receive the support they need to help them deliver their education and training programmes to learners.

For further information, please call our GCE line on 0844 576 0025, our GCSE team on 0844 576 0027, or visit our website at [www.edexcel.com](http://www.edexcel.com).

If you have any subject specific questions about the content of this Mark Scheme that require the help of a subject specialist, you may find our **Ask The Expert** email service helpful.

Ask The Expert can be accessed online at the following link:

<http://www.edexcel.com/Aboutus/contact-us/>


January 2010

Publications Code UA022965

All the material in this publication is copyright

© Edexcel Ltd 2010


January 2010  
6678 Mechanics M2  
Mark Scheme

Question Number	Scheme	Marks
Q1.	$\frac{dv}{dt} = 6t - 4$ $6t - 4 = 0 \Rightarrow t = \frac{2}{3}$ $s = \int 3t^2 - 4t + 3 dt = t^3 - 2t^2 + 3t (+c)$ $t = \frac{2}{3} \Rightarrow s = -\frac{16}{27} + 2 \text{ so distance is } \frac{38}{27} \text{ m}$	<p>M1 A1</p> <p>M1 A1</p> <p>M1 A1</p> <p>M1 A1</p> <p style="text-align: right;">[8]</p>
Q2.	<div style="text-align: center;">  </div> <p style="text-align: center;">CLM: <math>4mu - mu = 2mv_1 + mv_2</math></p> <p style="text-align: center;">i.e. <math>3u = 2v_1 + v_2</math></p> <p style="text-align: center;">NIL: <math>3eu = -v_1 + v_2</math></p> <p style="text-align: center;"><math>v_1 = u(1 - e)</math></p> <p style="text-align: center;"><math>v_2 = u(1 + 2e)</math></p>	<p>M1 A1</p> <p>M1 A1</p> <p>DM1 A1</p> <p>A1</p> <p style="text-align: right;">[7]</p>
Q3.	$\frac{1}{2} \times 0.5 \times 20^2 ; \quad 0.5g \times 10$ $10R = \frac{1}{2} \times 0.5 \times 20^2 - 0.5g \times 10$ $\Rightarrow R = 5.1$	<p>B1 B1</p> <p>M1 A1</p> <p>DM1 A1</p> <p style="text-align: right;">[6]</p>

Question Number	Scheme	Marks
Q4.	(i) $I \uparrow = 0.25 \times 40 \sin 60 = 5\sqrt{3}$ (8.66)      one component $I \leftarrow = 0.25(-20 + 30) = 2.5$ both $ I  = \sqrt{75 + 6.25} = 9.01$ (Ns)	M1 A1  M1 A1      (4)
	(ii) $\frac{\sin \theta}{40} = \frac{\sin 60^\circ}{\sqrt{1300}}$ $\theta = 106^\circ$ (3 s.f.)  or $\tan \theta = \pm \frac{5\sqrt{3}}{2.5}$ oee $\theta = 106^\circ$	M1 A1  M1 A1      (4)  [8]
	<i>Alternative to 4(i)</i> Use of $I = m(\mathbf{v} - \mathbf{u})$  $30^2 + 40^2 - 2 \times 30 \times 40 \cos 60^\circ$ (= 1300)  $I = 0.25\sqrt{1300} = 9.01$ N s (3 s.f.)	M1  M1 A1  A1
	<i>2nd Alternative to 4(i)</i> $\mathbf{u} = 30\mathbf{i}$ , $\mathbf{v} = 40 \cos 60\mathbf{i} + 40 \sin 60\mathbf{j} = 20\mathbf{i} + 20\sqrt{3}\mathbf{j}$ $I = \frac{1}{4}(-10\mathbf{i} + 20\sqrt{3}\mathbf{j}) = -2.5\mathbf{i} + 5\sqrt{3}\mathbf{j}$	M1 A1 etc

Question Number	Scheme	Marks
Q5.	<p>(a)</p> $\frac{490}{3.5} - R = 0$ $R = 140 \text{ N}$ <p>(b)</p> $\frac{24}{u} + 70g \cdot \frac{1}{14} - 40u = 0$ $40u^2 - 49u - 24 = 0$ $(5u - 8)(8u + 3) = 0$ $u = 1.6$	<p>B1 M1 A1</p> <p>A1 (4)</p> <p>B1</p> <p>M1 A2, 1, 0</p> <p>DM1</p> <p>DM1 A1 (7)</p> <p>[11]</p>
Q6.	$m(B) : R \times 4 \cos \alpha = F \times 4 \sin \alpha + 20g \times 2 \cos \alpha$ <p>Use of <math>F = \frac{1}{2}R</math></p> <p>Use of correct trig ratios</p> <p>R = 160N or 157N</p>	<p>M1 A2</p> <p>M1</p> <p>B1</p> <p>DM1 A1</p> <p>[7]</p>

Question Number	Scheme			Marks	
Q7.	(a)	<p>Rectangle</p> $24x$ $x$ $24x^2 - 4.5\pi \times \left(\frac{4 \times 3}{3\pi}\right) - 4.5\pi \times \left(\frac{4 \times 3}{3\pi}\right) = (24x + 9\pi)\bar{x}$ $\text{distance} =  \bar{x}  = \frac{4 2x^2 - 3 }{(8x + 3\pi)} \quad **$	<p>Semicircles</p> $4.5\pi \quad 4.5\pi$ $\frac{4 \times 3}{3\pi} \quad \frac{4 \times 3}{3\pi}$	<p>Template, <math>T</math></p> $24x + 9\pi$ $\bar{x}$	<p>B2</p> <p>B2</p> <p>M1 A1</p> <p>A1 (7)</p>
	(b)	<p>When <math>x = 2</math>,</p> $ \bar{x}  = \frac{20}{16 + 3\pi}$ $\tan \theta = \frac{6}{4 -  \bar{x} } = \frac{6}{4 - \frac{20}{16 + 3\pi}}$ $= \frac{48 + 9\pi}{22 + 6\pi} .$		<p>B1</p> <p>M1 A1</p> <p>A1 (4)</p> <p>[11]</p>	

Question Number	Scheme	Marks
Q8.	(a) $x = ut$	B1
	$y = cut - 4.9t^2$	M1 A1
	eliminating $t$ and simplifying to give $y = cx - \frac{4.9x^2}{u^2}$ **	DM1 A1 (5)
	(b)(i) $0 = cx - \frac{4.9x^2}{u^2}$	M1
	$0 = x(c - \frac{4.9x}{u^2}) \Rightarrow R = \frac{u^2c}{4.9} = 10c$	M1 A1
	(ii) When $x = 5c$ , $y = H$	M1
	$= 5c^2 - \frac{(5c)^2}{10} = 2.5c^2$	M1 A1 (6)
	(c) $\frac{dy}{dx} = c - \frac{9.8x}{u^2} = c - \frac{x}{5}$	M1 A1
	When $x = 0$ , $\frac{dy}{dx} = c$	B1
	So, $c - \frac{x}{5} = \frac{-1}{c}$	DM1 A1
$x = 5(c + \frac{1}{c})$	A1 (6)	
<p>Alternative to 8(c)</p>  <p><math>\tan \theta = \frac{u}{cu} = \frac{1}{c} = \frac{v}{u}</math></p> <p><math>\Rightarrow v = \frac{u}{c} = \frac{7}{c}</math></p> <p><math>v = u + at ; -\frac{7}{c} = 7c - 9.8t</math></p> <p><math>t = \frac{7}{9.8}(c + \frac{1}{c})</math></p> <p><math>x = ut = 7t ; x = 5(c + \frac{1}{c})</math></p>	<p>[17]</p> <p>B1</p> <p>M1 A1</p> <p>M1</p> <p>A1</p> <p>A1</p>	







Further copies of this publication are available from  
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467  
Fax 01623 450481

Email [publications@linneydirect.com](mailto:publications@linneydirect.com)

Order Code UA022965 January 2010

For more information on Edexcel qualifications, please visit [www.edexcel.com/quals](http://www.edexcel.com/quals)

Edexcel Limited. Registered in England and Wales no.4496750  
Registered Office: One90 High Holborn, London, WC1V 7BH