

# Mark Scheme (Results)

Summer 2012

GCE Mechanics M2  
(6678) Paper 1

## **Edexcel and BTEC Qualifications**

Edexcel and BTEC qualifications come from Pearson, the world's leading learning company. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information, please visit our website at [www.edexcel.com](http://www.edexcel.com).

Our website subject pages hold useful resources, support material and live feeds from our subject advisors giving you access to a portal of information. If you have any subject specific questions about this specification that require the help of a subject specialist, you may find our Ask The Expert email service helpful.

[www.edexcel.com/contactus](http://www.edexcel.com/contactus)

## **Pearson: helping people progress, everywhere**

Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: [www.pearson.com/uk](http://www.pearson.com/uk)

Summer 2012

Publications Code UA032678

All the material in this publication is copyright

© Pearson Education Ltd 2012

# Summer 2012 6678 Mechanics 2 Mark Scheme

## General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

# EDEXCEL GCE MATHEMATICS

## General Instructions for Marking

1. The total number of marks for the paper is 75.
2. The Edexcel Mathematics mark schemes use the following types of marks:
  - **M** marks: method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
  - **A** marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
  - **B** marks are unconditional accuracy marks (independent of M marks)
  - Marks should not be subdivided.
3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes and can be used if you are using the annotation facility on ePEN.

- bod – benefit of doubt
  - ft – follow through
  - the symbol  $\surd$  will be used for correct ft
  - cao – correct answer only
  - cso – correct solution only. There must be no errors in this part of the question to obtain this mark
  - isw – ignore subsequent working
  - awrt – answers which round to
  - SC: special case
  - oe – or equivalent (and appropriate)
  - dep – dependent
  - indep – independent
  - dp decimal places
  - sf significant figures
  - \* The answer is printed on the paper
  - $\square$  The second mark is dependent on gaining the first mark
4. All A marks are 'correct answer only' (cao.), unless shown, for example, as A1 ft to indicate that previous wrong working is to be followed through. After a misread however, the first 2 A or B marks affected are lost, and the subsequent A marks affected are treated as A ft; but manifestly absurd answers should never be awarded A marks.

## General Principles for Mechanics Marking

Usual rules for M marks: correct no. of terms; dim correct; all terms that need resolving (i.e. multiplied by cos or sin) are resolved.

Omission or extra  $g$  in a resolution is accuracy error not method error.

Omission of mass from a resolution is method error.

Omission of a length from a moments equation is a method error.

Omission of units or incorrect units is not (usually) counted as an accuracy error.

DM indicates a dependent method mark i.e. one that can only be awarded if a previous specified method mark has been awarded.

Any numerical answer which comes from use of  $g = 9.8$  should be given to 2 or 3 SF.

Use of  $g = 9.81$  should be penalised once per (complete) question.

N.B. Over-accuracy or under-accuracy of correct answers should only be penalised *ONCE* per complete question.

However, premature approximation should be penalised every time it occurs.

**MARKS MUST BE ENTERED IN THE SAME ORDER AS THEY APPEAR ON THE MARK SCHEME.**

In all cases, if the candidate clearly labels their working under a particular part of a question i.e. (a) or (b) or (c),.....then that working can only score marks for that part of the question.

Accept column vectors in all cases.

**Summer 2012**  
**6678 Mechanics M2**  
**Mark Scheme**


Question Number	Scheme	Marks	Notes
<b>1</b> <b>(a)</b>	$\mathbf{a} = \frac{d\mathbf{v}}{dt} = 6t\mathbf{i} + (4 - 2t)\mathbf{j}$ <p>When <math>t = 1</math>, <math>\mathbf{a} = 6\mathbf{i} + 2\mathbf{j}</math></p> $ \mathbf{a}  = \sqrt{6^2 + 2^2} = \sqrt{40} = 6.32 \text{ (m s}^{-2}\text{)}$	<p>M1</p> <p>A1</p> <p>DM1</p> <p>DM1</p> <p>A1</p>	<p>Differentiate <math>\mathbf{v}</math> to obtain <math>\mathbf{a}</math>.</p> <p>Accept column vector or <math>\mathbf{i}</math> and <math>\mathbf{j}</math> components dealt with separately.</p> <p>Substitute <math>t = 1</math> into their <math>\mathbf{a}</math>. Dependent on 1<sup>st</sup> M1</p> <p>Use of Pythagoras to find the magnitude of their <math>\mathbf{a}</math>.</p> <p>Allow with their t. Dependent on 1<sup>st</sup> M1</p> <p>Accept awrt 6.32, 6.3 or exact equivalents.</p>
	(5)	<p>M1</p> <p>A1</p> <p>DM1</p> <p>DM1</p> <p>A1</p>	
<b>(b)</b>	$\mathbf{r} = \int (3t^2 - 1)\mathbf{i} + (4t - t^2)\mathbf{j} dt$ $= (t^3 - t + C)\mathbf{i} + (2t^2 - \frac{1}{3}t^3 + D)\mathbf{j}$ <p><math>t = 0, \mathbf{r} = \mathbf{i} \Rightarrow C = 1, D = 0</math></p> <p>When <math>t = 3, \mathbf{r} = 25\mathbf{i} + 9\mathbf{j}</math> (m)</p>	<p>M1</p> <p>A1</p> <p>DM1</p> <p>DM1</p> <p>A1</p>	<p>Integrate <math>\mathbf{v}</math> to obtain <math>\mathbf{r}</math></p> <p>Condone <math>C, D</math> missing</p> <p>Use <math>t = 0, \mathbf{r} = \mathbf{i}</math> to find <math>C</math> &amp; <math>D</math></p> <p>Substitute <math>t = 3</math> with their <math>C</math> &amp; <math>D</math> to find <math>\mathbf{r}</math>. Dependent on both previous Ms.</p> <p>cao. Must be a vector.</p>
	(5)	(5)	
		<b>10</b>	

Question Number	Scheme	Marks	Notes
<p><b>2</b></p> <p><b>(a)</b></p> <p><b>(b)</b></p>	$3m \cdot 2u - 4mu = 3mv_1 + 4mv_2$ $e(2u + u) = -v_1 + v_2$ $\frac{u(2+9e)}{7} = v_2$ $v_1 = \frac{2u(1-6e)}{7}$ $v_1 < 0 \Rightarrow e > \frac{1}{6}$ $1 \geq e > \frac{1}{6}$	<p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>DM1</p> <p>A1</p> <p>(6)</p> <p>M1</p> <p>A1</p> <p>DM1</p> <p>A1</p> <p>B1</p> <p>(5)11</p>	<p>CLM. Need all terms. Condone sign slips.</p> <p>Correct but check their directions for <math>v_1</math> &amp; <math>v_2</math>.</p> <p>Impact law. Must be used the right way round, but condone sign slips.</p> <p>Directions of <math>v_1</math> &amp; <math>v_2</math> must be consistent between the two equations. (Ignore the diagram if necessary)</p> <p>Eliminate <math>v_1</math> to produce an equation in <math>v_2</math> only. Dependent on both previous M marks – must be using both equations.</p> <p>DO NOT accept the negative. The question asks for speed.</p> <p>Use the work from (a) or restart to find <math>v_1</math> or <math>\lambda v_1</math> for a constant <math>\lambda</math>. If using work from (a) this mark is dependent on the first 2 M marks.</p> <p>a.e.f. Correct for their direction. Allow for <math>\lambda v_1</math></p> <p>An appropriate inequality for their <math>v_1</math> (seen or implied) – requires previous M1 scored.</p> <p>Work on <math>v_1=0</math> scores M0 until the inequality is formed.</p> <p>Accept <math>\frac{2}{12}</math>. Answer must follow from correct work for <math>v_1</math></p> <p>For (their value) <math>&lt; e \leq 1</math></p> <p>SR: from <math>v_1 \leq 0</math> could score M1A0B1</p>

Question Number	Scheme	Marks	Notes
<p><b>3</b> <b>(a)</b></p> <p><math>M(A), F.4 \sin 40^\circ = 5g.2 \cos 25^\circ</math></p> <p><math>F = 35</math></p> <p><b>(b)</b></p> <p><math>F \cos 75^\circ \pm Y = 5g</math></p> <p><math>Y = 40 ;</math> UP</p>		<p>M1</p> <p>A1</p> <p>A1</p> <p>A1</p> <p>(4)</p> <p>M1</p> <p>A1</p> <p>A1</p> <p>A1</p> <p>(4)</p> <p><b>8</b></p>	<p>A complete method to find <math>F</math>, e.g. take moments about <math>A</math>. Condone sin/cos confusion. Requires correct ratio of lengths. Correct terms with at most one slip All correct 35 or 34.5 (&gt;3sf not acceptable due to use of 9.8, but only penalise once in a question)</p> <p>Resolve vertically. Need all three terms but condone sign errors. Must be attempting to work with their <math>75^\circ</math> or <math>15^\circ</math>. Correct equation (their <math>F</math>) 40 or 40.1 Apply ISW if the candidate goes on to find <math>R</math>. cso (the Q does specifically ask for the direction, so this must be clearly stated)</p>
<p><b>(b)</b></p>	<p>OR1: <math>4m \cos 25 \times Y</math> <math>= 5g \times 2m \cos 25 + F \cos 15 \times 4m \sin 25</math> etc.</p> <p>OR2: <math>R \cos \alpha = F \cos 40 + 5g \cos 65</math> <math>R \sin \alpha + F \sin 40 = 5g \cos 25</math> <math>R = 52.1, \alpha = 25.3^\circ</math> <math>Y = R \sin (25 + \alpha)</math> Etc.</p>	<p>M1</p> <p>A1</p> <p>M1A1</p>	<p>Taking moments about the point vertically below <math>B</math> and on the same horizontal level as <math>A</math>. (Their <math>F</math>)</p> <p>Resolve parallel &amp; perpendicular to the rod</p> <p>Solve for <math>R, \alpha</math></p> <p>Need a complete strategy to find <math>Y</math> for M1.</p>



Question Number	Scheme	Marks	Notes
<p><b>4</b></p> <p><b>(a)</b></p> <p><b>(b)</b></p>	$\frac{\pi(4a)^2}{4} \quad \frac{\pi(2a)^2}{2a} \quad (\frac{\pi(4a)^2 - \pi(2a)^2}{x})$ $(4 \times 4a) - (1 \times 2a) = 3 \bar{x}$ $\frac{14a}{3} = \bar{x} \quad *$ $OG = 4a \tan \alpha = \frac{10a}{3} \left( \Rightarrow PG = \frac{2a}{3} \right)$ $M(P), (m + km)g \cdot \frac{2a}{3} \cos \alpha = mg \cdot \frac{14a}{3} \cos \alpha$ $M(G) : km \times \frac{2}{3}a = m \times \left( \frac{10}{3}a + \frac{2}{3}a \right) = 4ma$ $M(O) : m(1+k) \times \frac{10}{3}a + m \times \frac{2}{3}a = km \times 4a$ $M(C) : \frac{12}{3}a \times (1+k)m = \frac{14}{3}a \times km$ $M(Q) : \frac{22}{3}a \times m(1+k) = \frac{10}{3}a \times m + 8a \times km$ $k = 6$	<p>B1</p> <p>B1</p> <p>M1</p> <p>A1</p> <p>(4)</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>A1</p> <p>(5)</p> <p><b>9</b></p>	<p>Correct mass ratios</p> <p>Distance of c of m from <math>P</math> (or from a point on <math>QP</math>).</p> <p>Moments about axis through <math>P</math>, or about a parallel axis then convert the answer to distance from <math>P</math>. Condone a sign slip. Answer given – check working carefully. Must reach positive answer legitimately.</p> <p>Vertical through <math>S</math> cuts <math>OP</math> at <math>G</math>. Use trig to find the position of <math>G</math> on <math>OP</math>.  <math>OG = \frac{10a}{3}</math>, <math>QG = \frac{22a}{3}</math> or <math>PG = \frac{2a}{3}</math> seen or implied</p> <p>Take moments about a point on <math>QP</math> – terms should be dimensionally consistent. Masses must be associated with the appropriate distances, which might be incorrectly evaluated or not yet found – e.g. accept with <math>QG</math>. Must have the right terms but condone trig confusion. Also condone absence of trig.</p> <p>cso (<math>C</math> is the position of the original centre of mass.)</p> <p>cso See next page for more alternatives.....</p>

<p><b>OR</b></p>	$(k+1)m \times PG = m \times \frac{14}{3}a$ $PG = \frac{14a}{3(k+1)}$ $\tan \alpha = \frac{OG}{4a} = \frac{4a - \frac{14a}{3(k+1)}}{4a} \left( = 1 - \frac{7}{6(k+1)} \right)$ $\frac{5}{6} = 1 - \frac{7}{6(k+1)} \quad k = 6$	<p>M1</p> <p>A1</p> <p>M1</p> <p>A1, A1</p>	<p>Moments about P</p> <p>Correct expression for PG</p> <p>Use of <math>\tan \alpha</math> in the correct triangle.</p> <p>Correct equation in <math>k</math>, correct solution</p>
<p><b>OR</b></p>	$\tan(\angle CSO) = \tan \beta = \frac{2a}{3} = \frac{1}{6}$  $km \cdot \sqrt{32}a \cdot \sin(45 - \alpha) = m \cdot \sqrt{16 \frac{4}{9}}a \cdot \sin(\alpha + \beta)$ $k \cdot \sqrt{32} \cdot \left( \frac{6-5}{\sqrt{2} \cdot \sqrt{61}} \right) = \frac{\sqrt{148}}{3} \cdot \frac{6 \times 5 + 1 \times 6}{\sqrt{37} \cdot \sqrt{61}}$ $4k = \frac{2}{3} \times 36, \quad k = 6$	<p>M1</p>   <p>M1A1</p> <p>A1</p> <p>A1</p>	 <p>Moments about S</p> <p>Do not expect accurate working</p> <p>Final answer 6.0</p>



Question Number	Scheme	Marks	Notes	
<b>6</b>	<b>(a)</b> $F = \frac{60000}{10} = 6000$ $F - 1200g \sin \alpha - 400g \sin \alpha - 1000 - 200 = 1600a$ $a = 2.3 \text{ (m s}^{-2}\text{)}$	B1	Correct application of $P = Fv$ seen or implied.	
		M1	Use of $F = ma$ parallel to the slope for the car and trailer.	
		A1	Must have all the terms, but condone sign errors.	
		A1	At most one error (with $F$ or their $F$ )	
		A1	Correct equation (with $F$ or their $F$ ) only	
	<b>(b)</b> $T - 400g \sin \alpha - 200 = 400 \times 2.3$ $T = 1400$	(5)	M1	Use of $F = ma$ parallel to the slope for the trailer
		A1 ft	A1 ft	At most one error (their $a$ )
		A1	A1	All correct (their $a$ ) only
	<b>OR</b> $6000 - 1200g \sin \alpha - 1000 - T = 1200 \times 2.3$ $T = 1400$	(4)	M1	Use of $F = ma$ parallel to the slope for the car
		A1 ft	A1 ft	At most one error (their $a$ )
A1		A1	All correct (their $a$ ) only	
<b>OR (a)</b> $F = 6000$ $T - 400g \sin \alpha - 200 = 400 \times a$ $6000 - 1200g \sin \alpha - 1000 - T = 1200 \times a$ $6000 - 1600g \sin \alpha - 1200 = 1600a$ $a = 2.3 \text{ (m s}^{-2}\text{)}$	(4)	B1	Simultaneous equations in $T$ and $a$	
		M1A1A1	Add to eliminate $T$	
	<b>(b)</b> $-800a = 2T + 800g \sin \alpha + 800 - 6000$ $2T = 5200 - 800g \sin \alpha - 800 \times 2.3$ $T = 1400$	A1	M1A1A1	Subtract and / or substitute to eliminate $a$
		A1	A1	
		A1	A1	

<p>(c)</p>	$200d = \frac{1}{2} 400.12^2 - 400gd \sin \alpha$ $d = 60 \text{ (m)}$	<p>M1</p> <p>A1</p> <p>A1</p> <p>DM1</p> <p>A1</p> <p>(5)</p> <p><b>14</b></p>	<p>Use of work-energy. Must have all three terms. Do not accept duplication of terms, but condone sign errors.</p> <p>Equation in only one unknown, but could be vertical distance.</p> <p>At most one error in the equation</p> <p>All correct in one unknown</p> <p>Solve for <math>d</math> – dependent on M for work-energy equation. only</p> <p>For vertical distance <math>\left( = \frac{60}{14} = 4.29 \right)</math> allow 3/5</p>
------------	---	--	--

Question Number	Scheme	Marks	
<p><b>7 (a)</b></p> <p>OR</p> <p><b>(b)</b></p>	$0^2 = u_v^2 - 2 \times 9.8 \times 10$ $u_v = 14 \quad *$	M1 A1 A1 (3)	Complete method using <i>suvat</i> to form an equation in $u_v$ . Correct equation e.g. $0 = u^2 - 20g$ *Answer given* requires equation and working, including 196, seen.
	conservation of energy: $\frac{1}{2} m (u_h^2 + u_v^2) = mg \times 10 + \frac{1}{2} m u_h^2, \frac{1}{2} u_v^2 = 98$ $u_v = 14 \quad *$	M1 A1 A1 (3)	Initial KE = gain in GPE + final KE Correct equation *Answer given*
	$(\uparrow), -52.5 = 14t - \frac{1}{2} g t^2$	M1 A1 A1	Use the vertical distance travelled to find the <b>total</b> time taken. At most one error Correct equation
	$49t^2 - 140t - 525 = 0$ $(t - 5)(49t + 105) = 0 \quad t = 5$	DM1 A1	Solve for $t$ . Dependent on the preceding M mark only
	$(\rightarrow), 50 = 5u_H$ $u_H = 10$	M1 A1	Use their time of flight to form an equation in $u_H$ only
	$u = \sqrt{10^2 + 14^2}$	M1	Use of Pythagoras with two non-zero components, or solution of a pair of simultaneous equations in $u$ and $\alpha$ .
	$= \sqrt{296}; 17.2 \text{ m s}^{-1}$	A1	17.2 or 17 (method involves use of $g = 9.8$ so an exact surd answer is not acceptable)
			(9) See next page for an alternative route to $u$ , and (c).

<p><b>OR</b></p> $50 = u \cos \alpha t \quad \text{or} \quad 50 = u_H t$ $49 \left( \frac{50}{u_H} \right)^2 - 140 \left( \frac{50}{u_H} \right) - 525 = 0$ $525(u_H)^2 + 140(u_H) - 122500 = 0$ <p>Solve for <math>u_H</math></p> $u_H = 10$ <p>etc.</p>		<p>M1</p> <p>A1</p> <p>DM1</p> <p>A1</p>	<p>First 3 marks for the quadratic as above. Used in their quadratic</p> <p>Correct quadratic in <math>u_H</math></p> <p>Dependent on the M mark for setting up the initial quadratic equation in t. only Complete as above.</p>
<p>(c)</p> $\tan OBA = \frac{52.5}{50} = 1.05$ $v_V = 1.05 \times 10 = 10.5$ <p>(<math>\uparrow</math>), <math>-10.5 = 14 - gt</math></p> $t = 2.5$		<p>B1</p> <p>M1</p> <p>DM1</p> <p>A1</p> <p>A1</p> <p>(5)</p> <p><b>17</b></p>	<p>Correct direction o.e. (accept reciprocal)</p> <p>Use trig. with their <math>u_H</math> and correct interpretation of direction to find the vertical component of speed. Working with distances is M0. (condone <math>10 \div 1.05</math>) Use suvat to form an equation in t. Dependent on the preceding M. Correct equation for their <math>u_H</math>. For incorrect direction give A0 here. only</p>

Further copies of this publication are available from  
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467

Fax 01623 450481

Email [publication.orders@edexcel.com](mailto:publication.orders@edexcel.com)

Order Code UA032678 Summer 2012

For more information on Edexcel qualifications, please visit our website  
[www.edexcel.com](http://www.edexcel.com)

Pearson Education Limited. Registered company number 872828  
with its registered office at Edinburgh Gate, Harlow, Essex CM20 2JE

Ofqual  




Llywodraeth Cynulliad Cymru  
Welsh Assembly Government

