



1.




Figure 1

Two smooth uniform spheres *A* and *B* have masses  $2m$  kg and  $3m$  kg respectively and equal radii. The spheres are moving on a smooth horizontal surface. Initially, sphere *A* has velocity  $(3\mathbf{i} - 4\mathbf{j}) \text{ m s}^{-1}$  and sphere *B* has velocity  $(2\mathbf{i} + 3\mathbf{j}) \text{ m s}^{-1}$ . When the spheres collide, the line joining their centres is parallel to  $\mathbf{j}$ , as shown in Figure 1. The coefficient of restitution between the spheres is  $\frac{3}{7}$ . Find, in terms of  $m$ , the total kinetic energy lost in the collision.

(10)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---





2.




Figure 2

Figure 2 represents part of the smooth rectangular floor of a sports hall. A ball is at  $B$ , 4 m from one wall of the hall and 5 m from an adjacent wall. These two walls are smooth and meet at the corner  $C$ . The ball is kicked so that it travels along the floor, bounces off the first wall at the point  $X$  and hits the second wall at the point  $Y$ . The point  $Y$  is 7.5 m from the corner  $C$ .

The coefficient of restitution between the ball and the first wall is  $\frac{3}{4}$ .

Modelling the ball as a particle, find the distance  $CX$ .

(9)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---





3. [In this question the unit vectors  $\mathbf{i}$  and  $\mathbf{j}$  are due east and due north respectively.]

A coastguard patrol boat  $C$  is moving with constant velocity  $(8\mathbf{i} + u\mathbf{j}) \text{ km h}^{-1}$ . Another ship  $S$  is moving with constant velocity  $(12\mathbf{i} + 16\mathbf{j}) \text{ km h}^{-1}$ .

(a) Find, in terms of  $u$ , the velocity of  $C$  relative to  $S$ . (2)

At noon,  $S$  is 10 km due west of  $C$ .  
If  $C$  is to intercept  $S$ ,

(b) (i) find the value of  $u$ .  
(ii) Using this value of  $u$ , find the time at which  $C$  would intercept  $S$ . (4)

If instead, at noon,  $C$  is moving with velocity  $(8\mathbf{i} + 8\mathbf{j}) \text{ km h}^{-1}$  and continues at this constant velocity,

(c) find the distance of closest approach of  $C$  to  $S$ . (5)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



Question 3 continued

Lined writing area for the answer to Question 3 continued.



P 3 5 4 1 5 A 0 7 2 4







4. A hiker walking due east at a steady speed of  $5 \text{ km h}^{-1}$  notices that the wind appears to come from a direction with bearing 050. At the same time, another hiker moving on a bearing of 320, and also walking at  $5 \text{ km h}^{-1}$ , notices that the wind appears to come from due north.

Find

(a) the direction from which the wind is blowing, (3)

(b) the wind speed. (4)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---





5. A particle  $Q$  of mass 6 kg is moving along the  $x$ -axis. At time  $t$  seconds the displacement of  $Q$  from the origin  $O$  is  $x$  metres and the speed of  $Q$  is  $v$  m s<sup>-1</sup>. The particle moves under the action of a retarding force of magnitude  $(a + bv^2)$  N, where  $a$  and  $b$  are positive constants. At time  $t = 0$ ,  $Q$  is at  $O$  and moving with speed  $U$  m s<sup>-1</sup> in the positive  $x$ -direction. The particle  $Q$  comes to instantaneous rest at the point  $X$ .

(a) Show that the distance  $OX$  is

$$\frac{3}{b} \ln \left( 1 + \frac{bU^2}{a} \right) \text{ m} \quad (6)$$

Given that  $a = 12$  and  $b = 3$ ,

(b) find, in terms of  $U$ , the time taken to move from  $O$  to  $X$ . (5)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---







**Question 5 continued**

Ruled area for writing the answer to Question 5 continued.

**Q5**

--	--

**(Total 11 marks)**







**Question 6 continued**

Lined area for writing the answer to Question 6.





**Question 6 continued**

Lined writing area consisting of multiple horizontal lines for student response.


**(Total 13 marks)**

**Q6**

--	--



7.



**Figure 3**

Figure 3 shows a framework  $ABC$ , consisting of two uniform rods rigidly joined together at  $B$  so that  $\angle ABC = 90^\circ$ . The rod  $AB$  has length  $2a$  and mass  $4m$ , and the rod  $BC$  has length  $a$  and mass  $2m$ . The framework is smoothly hinged at  $A$  to a fixed point, so that the framework can rotate in a fixed vertical plane. One end of a light elastic string, of natural length  $2a$  and modulus of elasticity  $3mg$ , is attached to  $A$ . The string passes through a small smooth ring  $R$  fixed at a distance  $2a$  from  $A$ , on the same horizontal level as  $A$  and in the same vertical plane as the framework. The other end of the string is attached to  $B$ . The angle  $ARB$  is  $\theta$ , where  $0 < \theta < \frac{\pi}{2}$ .

(a) Show that the potential energy  $V$  of the system is given by

$$V = 8amg \sin 2\theta + 5amg \cos 2\theta + \text{constant} \quad (7)$$

(b) Find the value of  $\theta$  for which the system is in equilibrium. (4)

(c) Determine the stability of this position of equilibrium. (3)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---







Leave  
blank

**Question 7 continued**

Handwriting lines for the answer to Question 7.

**Q7**

**(Total 14 marks)**

**TOTAL FOR PAPER: 75 MARKS**

**END**

