

Mark Scheme (Results)

Summer 2012

GCE Statistics S4
(6686) Paper 1

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications come from Pearson, the world's leading learning company. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information, please visit our website at www.edexcel.com.

Our website subject pages hold useful resources, support material and live feeds from our subject advisors giving you access to a portal of information. If you have any subject specific questions about this specification that require the help of a subject specialist, you may find our Ask The Expert email service helpful.

www.edexcel.com/contactus

Pearson: helping people progress, everywhere

Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Summer 2012

Publications Code UA033143

All the material in this publication is copyright

© Pearson Education Ltd 201

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

June 2012
6686 Statistics S4
Mark Scheme

Question Number	Scheme	Marks
1 (a)	<p>$H_0: \mu_d = 0, H_1: \mu_d > 0$ (or $H_1: \mu_d < 0$)</p> <p>where μ_d is the (population) mean difference :- BP sitting down – BP standing. (BP standing – BP sitting down)</p> <p>Assume the differences are normally distributed</p>	<p>B1</p> <p>B1</p> <p>(2)</p>
1 (b)	<p>$d: 4, -1, 6, 6, 3, -2, 9, -1, 4, 7, -11, 7$</p> <p>$(\Sigma d = 31, \Sigma d^2 = 419) \bar{d} = \pm 2.5833; sd = 5.55073. (or Var = 30.8106)$</p> <p>$t = \frac{\pm 2.5833\sqrt{12}}{5.55073} = \pm 1.612\dots$ Formula and substitution, 1.61</p> <p>Critical value $t_{11}(1\%) = 2.718(1 \text{ tail})$</p> <p>Not significant. Insufficient evidence to support that the blood pressure of a person sitting down is more than the blood pressure of a person after standing up.</p>	<p>M1</p> <p>A1; A1</p> <p>M1, A1</p> <p>B1</p> <p>A1 ft</p> <p>(7)</p> <p>Total 9 marks</p>
1a	<p>Notes</p> <p>B1 both hypotheses.</p> <p>B1 must be differences</p>	
1b	<p>M1 at least 2 correct or may be implied by correct Σd or Σd^2 or \bar{d} or sd or var or implied by correct t value</p> <p>A1 correct \bar{d} awrt ± 2.58- may be implied by correct t value</p> <p>A1 correct sd awrt 5.55 or var awrt 30.8 - may be implied by correct t value</p> <p>M1 $\frac{\pm \text{their } \bar{d} \sqrt{12}}{\text{their sd}}$</p> <p>A1 awrt 1.61</p> <p>B1 CV</p> <p>A1ft follow through their t value – need context of blood pressure and sitting and standing</p>	

Question Number	Scheme	Marks
2 (a)	$S_F^2 = \frac{1}{5} \{2308.01 - 6 \times 19.6^2\} = 0.61$ $S_M^2 = \frac{1}{11} \{2262.57 - 12 \times 13.7^2\} = 0.93545..$ <p>$H_0: \mu_F = \mu_M + 5; H_1: \mu_F \neq \mu_M + 5$ both</p> <p>CR: $t_{16}(0.025) > 2.120$ 2.12</p> $S_p^2 = \frac{5 \times 0.61 + 11 \times 0.93545...}{16} = 0.83375$ $t = \frac{19.6 - 13.7 - 5}{\sqrt{0.83375 \left(\frac{1}{6} + \frac{1}{12}\right)}} = 1.971$ <p>Since 1.971 is not in the critical region we accept H_0 and conclude that the mean shell length of female turtles does exceed the shell length of male turtles by 5cm.(or Biologists claim is correct)</p>	<p>B1</p> <p>B1</p> <p>B1</p> <p>B1</p> <p>M1 A1</p> <p>M1 A1ftA1</p> <p>A1 ft</p> <p>(10)</p> <p>B1 M1</p> <p>A1cso</p> <p>M1</p> <p>M1</p> <p>A1 awrt 0.44</p> <p>(6)</p> <p>Total 16 marks</p>
(b)(i)	$-1.96 < \frac{\bar{X}_F - \bar{X}_M - 5}{\sqrt{\left(\frac{0.9}{6} + \frac{0.9}{12}\right)}} < 1.96$ $4.07 < \bar{X}_F - \bar{X}_M < 5.93$	
(ii)	<p>P(Type II error) = $P(4.07 < \bar{X}_F - \bar{X}_M < 5.93 \mid N(6, 0.225))$</p> $= P\left(\frac{4.07 - 6}{\sqrt{0.225}} < z < \frac{5.93 - 6}{\sqrt{0.225}}\right)$ $= 0.44$ <p>awrt 0.44</p>	
2(a)	<p>B1 – awrt 0.61</p> <p>B1 – awrt 0.935</p> <p>Both may be implied by correct t value or S_p</p> <p>B1 allow rearrangements eg $\mu_F - \mu_M = 5$. If M and F not used then they must make clear what each letter is.</p> <p>B1 CV (if using one tail test allow 1.746)</p> <p>M1 $\frac{5 \times \text{their } 0.61 + 11 \times \text{their } 0.93545...}{16}$</p> <p>A1 awrt 0.834</p> <p>M1 $\pm \left(\frac{19.6 - 13.7 - 5}{\sqrt{p \left(\frac{1}{6} + \frac{1}{12}\right)}}\right)$ where p is either their 0.61 or 0.94 or their S_p^2 (awrt 0.834) (Allow $13.7 - 19.6 - 5$)</p> <p>A1 ft their S_p^2</p> <p>A1 awrt 1.97</p>	
(b)	<p>B1 1.96</p> <p>M1 must use z value</p>	
(c)	<p>M1 writing or using $N(6, 0.225)$</p> <p>M1 finding correct area and standardising (must use 6 but allow use of 0.9 and $(0.9/18)$ for var)</p>	

Question Number	Scheme	Marks
3.	<p data-bbox="316 241 699 286">$H_0: \sigma_A^2 = \sigma_B^2; H_1: \sigma_A^2 \neq \sigma_B^2$</p> <p data-bbox="320 309 815 389">$S_A^2 / S_B^2 = \frac{225}{36} = 6.25 \left(\frac{36}{225} = 0.16 \right)$</p> <p data-bbox="309 474 719 560">CR: $F_{10,8} > 3.35 \left(\frac{1}{F_{10,8}} = 0.299 \right)$</p> <p data-bbox="220 586 1329 658">Since 6.25 is in the critical region we can assume that the lengths of paving slabs sold by the builders merchant differ in variability.</p> <p data-bbox="220 824 1090 860">B1 both correct. Must use σ. May use different notation to A and B</p> <p data-bbox="220 882 660 958">M1 $\frac{225}{36}$ or $\frac{36}{225}$ allow $\frac{15}{6}$ or $\frac{6}{15}$</p> <p data-bbox="220 981 507 1016">A1 either 6.25 or 0.16</p> <p data-bbox="220 1039 643 1075">B1 CR must match their method</p> <p data-bbox="220 1097 775 1133">A1 context must include "lengths of slabs"</p>	<p data-bbox="1361 241 1406 277">B1</p> <p data-bbox="1361 309 1445 344">M1A1</p> <p data-bbox="1361 474 1406 510">B1</p> <p data-bbox="1361 586 1426 622">A1ft</p> <p data-bbox="1481 658 1522 694">(5)</p> <p data-bbox="1406 712 1481 748">Total</p> <p data-bbox="1406 766 1522 801">5 marks</p>

Question Number	Scheme	Marks
<p>4</p> <p>(a)</p> <p>(i)</p> <p>(ii)</p> <p>(b)</p>	<p>$\bar{x} = 4.9$</p> <p>$s = \sqrt{0.191..} \quad (0.437...)$</p> <p>(NB: $\Sigma x = 49$; $\Sigma x^2 = 241.82$)</p> <p>95% confidence interval is given by</p> <p>$4.9 \pm 2.262 \times \sqrt{\frac{0.191..}{10}}$</p> <p>i.e: (4.587..., 5.212 ...)</p> <p>95% confidence interval is given by</p> <p>$\frac{9 \times 0.437...^2}{19.023} < \sigma^2 < \frac{9 \times 0.437...^2}{2.7}$ use of $\frac{(n-1)s^2}{\chi_{n-1}^2}$</p> <p>i.e; (0.0904, 0.63704)</p> <p>5 lies inside the confidence interval</p> <p>0.49(0.7²) lies inside the confidence interval</p> <p>Yes it does meet the time requirement</p>	<p>B1</p> <p>B1</p> <p>M1A1ft B1</p> <p>A1 A1</p> <p>M1B1B1A1</p> <p>A1 A1</p> <p>(13)</p> <p>B1ft</p> <p>B1ft</p> <p>B1 ft</p> <p>(3)</p> <p>Total</p> <p>16 marks</p>

Question Number	Scheme	Marks
(a)	<p>B1 B1 may be implied by correct a correct answer to (i) or (ii)</p> <p>(i) M1 - “their 4.9” \pm t value $\times \sqrt{\frac{\text{their } 0.191..}{10}}$</p> <p>A1ft - “their 4.9” \pm 2.262 $\times \sqrt{\frac{\text{their } 0.191..}{10}}$</p> <p>B1 2.262</p> <p>A1 either correct to 3sf or better or both correct to 2sf or better</p> <p>A1 both correct to 3sf or better</p> <p>(ii) M1 – writing and attempting to use $\frac{(n-1)s^2}{\chi_{n-1}^2}$ or may be implied by correct formula used with their 0.437</p> <p>B1 19.023</p> <p>B1 2.7</p> <p>A1ft follow through their 0.437 and two chi squared values</p> <p>A1 either correct to 2sf or better</p> <p>A1 awrt (0.09, 0.637)</p> <p>(b) For the second B1. If both 0.7 and 0.49 lie in interval they must state variance = 0.49 or the interval for standard deviation.</p> <p>For the third B1 their must not be two conflicting conclusions unless they give just one overall as well.</p>	

Question Number	Scheme	Marks
5.(a)	$H_0: \sigma^2 = 36; H_1: \sigma^2 > 36$ $v = 24, X_{24}^2(0.05) = 36.415$ $\frac{(n-1)S^2}{\sigma^2} = \frac{24 \times 55}{36} = 36.67$ <p>Since $36.67 > 36.415$ there is sufficient evidence to reject H_0. There is evidence to suggest that the variance is greater than 36.</p>	B1 B1 M1 A1 A1 ft A1 ft (6)
(b)	$H_0: \mu = 450 \quad H_1: \mu > 450$ $t_{24} = 1.711$ $t = \pm \frac{455 - 450}{\sqrt{\frac{55}{25}}} = \pm 3.37\dots$ <p>Significant; The <u>mean weight</u> of chocolates is <u>greater than 450</u>. Or <u>μ is more than 450</u></p>	B1 B1 M1 A1 A1ft; A1ft (6)
(c)	<p>The <u>weights</u> are normally distributed</p>	B1 (1) Total 13 marks
(a)	<p>Notes</p> <p>B1 both correct. Also allow $H_0: \sigma = 6; H_1: \sigma > 6$ B1 36.415 M1 use of $\frac{(n-1)S^2}{\sigma^2}$ A1 awrt 36.7</p>	
(b)	$M1 \pm \frac{455 - 450}{\sqrt{\frac{55}{25}}}$ <p>A1 awrt 3.4</p> <p>A1ft any statement – no conflicting A1ft contextual statement must include “weight of chocolate” and is “greater than 50”</p>	

Question Number	Scheme	Marks
6(a)(i)	$E(\hat{p}_1) = E\left(\frac{X}{n}\right)$ $= \frac{1}{n} E(X)$ $= \frac{1}{n} \times np$ $= p \quad \text{unbiased}$	M1 A1cso
(ii)	$\text{Var}(\hat{p}_1) = \text{Var}\left(\frac{X}{n}\right)$ $= \frac{1}{n^2} \text{Var}(X)$ $= \frac{1}{n^2} \times np(1-p)$ $= \frac{p(1-p)}{n}$	M1 A1
b(i)	$E(\hat{p}_3) = 3a E(\hat{p}_1) + 2a E(\hat{p}_2)$ $= 3ap + 2ap$ $= 5ap$ $5ap = p$ $a = \frac{1}{5}$	M1 M1 A1
(ii)	$\text{Var}(\hat{p}_3) = \frac{9}{25} \text{Var}(\hat{p}_1) + \frac{4}{25} \text{Var}(\hat{p}_2)$ $= \frac{9p(1-p)}{25n} + \frac{4p(1-p)}{25m}$ $= \frac{p(1-p)}{25} \left(\frac{9}{n} + \frac{4}{m}\right)$	M1 M1d A1
(c)	$\frac{p(1-p)}{25} \left(\frac{9}{n} + \frac{4}{m}\right) < \frac{p(1-p)}{n}$ $9m + 4n < 25m$ $4n < 16m$ $\frac{n}{m} < 4$ $\frac{p(1-p)}{25} \left(\frac{9}{n} + \frac{4}{m}\right) < \frac{p(1-p)}{m}$ $9m + 4n < 25n.$	M1 M1

(4)

(6)

Question Number	Scheme	Marks
(d)	$9m < 21n$ $\frac{9}{21} < \frac{n}{m} \text{ or } \frac{3}{7} < \frac{n}{m}$ $\frac{3}{7} < \frac{n}{m} < 4$ <p> $\text{Var}(\hat{p}_1) = 0.05 p(1 - p)$ $\text{Var}(\hat{p}_2) = 0.0167 p(1 - p)$ $\text{Var}(\hat{p}_3) = 0.0207 p(1 - p)$ </p> <p>Or since $\frac{1}{3}$ is not in the range $\frac{9}{21} < \frac{n}{m} < 4$ $\text{Var}(\hat{p}_3)$ is not the smallest variance.</p> <p> $\text{Var}(\hat{p}_1) = 0.05 p(1 - p)$ $\text{Var}(\hat{p}_2) = 0.0167 p(1 - p)$ </p> <p>Therefore \hat{p}_2; is the best estimator as it has the smallest variance</p> <p>Notes</p> <p>(a) (i) M1 either $\frac{1}{n} E(X)$ or $\frac{1}{n} \times np$</p> <p>A1 cso</p> <p>(ii) M1 either $\frac{1}{n^2} \text{Var}(X)$ or $\frac{1}{n^2} \times np(1 - p)$</p> <p>A1 cso</p> <p>(b) (i) M1 For either $3a E(\hat{p}_1) + 2a E(\hat{p}_2)$ or $3ap + 2ap$</p> <p>M1 Putting their $E(\hat{p}_3) = p$</p> <p>(ii) M1 for $\frac{9}{25} \text{Var}(\hat{p}_1) + \frac{4}{25} \text{Var}(\hat{p}_2)$</p> <p>M1d for substituting (aii) for $\text{Var}(\hat{p}_1)$ and (aii) with m instead of n for $\text{Var}(\hat{p}_2)$</p> <p>A1 cso</p> <p>(c) M1 Putting $\text{Var}(\hat{p}_3) < \text{their } \text{Var}(\hat{p}_1)$ leading to an inequality of the form $\frac{n}{m} < a$ or $\frac{n}{m} > a$ where a is a constant.</p> <p>M1 Putting $\text{Var}(\hat{p}_3) < \text{their } \text{Var}(\hat{p}_2)$ leading to an inequality of the form $\frac{n}{m} > a$ or</p>	<p>A1</p> <p>(3)</p> <p>M1</p> <p>A1ft; A1ft</p> <p>(3)</p> <p>Total 16 marks</p>

$$\frac{n}{m} < a \text{ where } a \text{ is a constant.}$$

(d)

1/3 is not in their range in part(c)

M1 attempt to find all 3 variances or eliminating $\text{Var}(\hat{p}_3)$ with reason and finding the other 2 variances.

A1ft correct estimator chosen.

A1ft correct supporting reason from correct working for their var formulae

SC if 1/3 is in their range in part(c) they may get

B1 for stating \hat{p}_3

B1 dependent on the previous B being awarded- stating smallest variance award first two marks on open.

Further copies of this publication are available from
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467
Fax 01623 450481
Email publication.orders@edexcel.com
Order Code UA031443 Summer 2012

For more information on Edexcel qualifications, please visit our website
www.edexcel.com

Pearson Education Limited. Registered company number 872828
with its registered office at Edinburgh Gate, Harlow, Essex CM20 2JE

Ofqual
.....



Llywodraeth Cynulliad Cymru
Welsh Assembly Government

