

Question 1 continued

Leave
blank

Q1

(Total 5 marks)



N 2 6 1 1 0 A 0 3 2 4

Leave
blank

5.

The line l_1 has equation $\mathbf{r} = \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix} + \lambda \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}$.

The line l_2 has equation $\mathbf{r} = \begin{pmatrix} 1 \\ 3 \\ 6 \end{pmatrix} + \mu \begin{pmatrix} 2 \\ 1 \\ -1 \end{pmatrix}$.

(a) Show that l_1 and l_2 do not meet.

(4)

The point A is on l_1 where $\lambda = 1$, and the point B is on l_2 where $\mu = 2$.

(b) Find the cosine of the acute angle between AB and l_1 .

(6)



Question 5 continued

Leave
blank

Lined area for writing the answer to Question 5.

(Total 10 marks)

Q5



N 2 6 1 1 0 A 0 1 1 2 4

Question 6 continued

Lined writing area for the answer to Question 6.

Leave blank

Q6

(Total 12 marks)



7.




Figure 1

Figure 1 shows part of the curve with equation $y = \sqrt{(\tan x)}$. The finite region R , which is bounded by the curve, the x -axis and the line $x = \frac{\pi}{4}$, is shown shaded in Figure 1.

- (a) Given that $y = \sqrt{(\tan x)}$, complete the table with the values of y corresponding to $x = \frac{\pi}{16}$, $\frac{\pi}{8}$ and $\frac{3\pi}{16}$, giving your answers to 5 decimal places.

x	0	$\frac{\pi}{16}$	$\frac{\pi}{8}$	$\frac{3\pi}{16}$	$\frac{\pi}{4}$
y	0				1

(3)

- (b) Use the trapezium rule with all the values of y in the completed table to obtain an estimate for the area of the shaded region R , giving your answer to 4 decimal places.

(4)

The region R is rotated through 2π radians around the x -axis to generate a solid of revolution.

- (c) Use integration to find an exact value for the volume of the solid generated.

(4)





Question 7 continued

Leave
blank

Lined area for writing the answer to Question 7.



N 2 6 1 1 0 A 0 1 7 2 4



8. A population growth is modelled by the differential equation

$$\frac{dP}{dt} = kP,$$

where P is the population, t is the time measured in days and k is a positive constant.

Given that the initial population is P_0 ,

- (a) solve the differential equation, giving P in terms of P_0 , k and t . (4)

Given also that $k = 2.5$,

- (b) find the time taken, to the nearest minute, for the population to reach $2P_0$. (3)

In an improved model the differential equation is given as

$$\frac{dP}{dt} = \lambda P \cos \lambda t,$$

where P is the population, t is the time measured in days and λ is a positive constant.

Given, again, that the initial population is P_0 and that time is measured in days,

- (c) solve the second differential equation, giving P in terms of P_0 , λ and t . (4)

Given also that $\lambda = 2.5$,

- (d) find the time taken, to the nearest minute, for the population to reach $2P_0$ for the first time, using the improved model. (3)





Question 8 continued

Leave
blank

Lined area for writing the answer to Question 8.





Question 8 continued

Leave
blank

Lined area for writing answers.

Q8

(Total 14 marks)

TOTAL FOR PAPER: 75 MARKS

END



N 2 6 1 1 0 A 0 2 3 2 4

BLANK PAGE

