

Please check the examination details below before entering your candidate information

Candidate surname		Other names	
Centre Number		Candidate Number	

Pearson Edexcel Level 3 GCE

Monday 16 June 2025

Morning (Time: 1 hour 45 minutes)

Paper reference **9CH0/02**

Chemistry
Advanced
PAPER 2: Advanced Organic and Physical Chemistry

You must have:
 Scientific calculator, Data Booklet, ruler

Total Marks

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided
 – *there may be more space than you need.*

Information

- The total mark for this paper is 90.
- The marks for **each** question are shown in brackets
 – *use this as a guide as to how much time to spend on each question.*
- For the question marked with an **asterisk** (*), marks will be awarded for your ability to structure your answer logically, showing how the points that you make are related or follow on from each other where appropriate.
- A Periodic Table is printed on the back cover of this paper.

Advice

- Read each question carefully before you start to answer it.
- Show all your working in calculations and include units where appropriate.
- Check your answers if you have time at the end.

Turn over ►

P77833A

©2025 Pearson Education Ltd.
 Y:1/1/1/




Pearson

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

BLANK PAGE



Answer ALL questions.

Some questions must be answered with a cross in a box ☐. If you change your mind about an answer, put a line through the box ☒ and then mark your new answer with a cross ☐.

1 This question is about mass spectrometry.

(a) Which molecule would **not** give a molecular ion peak with $m/z = 74$?

(1)

- ☐ **A** $\text{C}_2\text{H}_5\text{COOH}$
☐ **B** CHOCOOH
☐ **C** $\text{CH}_2(\text{OH})\text{COOH}$
☐ **D** $\text{CH}_3\text{COOCH}_3$

(b) The fragmentation pattern in a mass spectrum can be used to distinguish between structural isomers.

Which m/z peak would be present in the mass spectrum of propanal, $\text{CH}_3\text{CH}_2\text{CHO}$, but **not** in the mass spectrum of propanone, CH_3COCH_3 ?

(1)

- ☐ **A** 15
☐ **B** 29
☐ **C** 43
☐ **D** 59

(Total for Question 1 = 2 marks)

2 Intermolecular forces affect the physical properties of compounds.

- (a) Which of these isomers, with the formula C_8H_{18} , will have the **lowest** boiling temperature?

(1)

- ☐ **A** $(CH_3)_2CHCH_2C(CH_3)_3$
- ☐ **B** $(CH_3)_2CHCH_2CH(CH_3)CH_2CH_3$
- ☐ **C** $CH_3CH_2CH_2C(CH_3)_2CH_2CH_3$
- ☐ **D** $CH_3CH_2CH_2CH_2CH_2CH_2CH_2CH_3$

- (b) Pentan-1-ol and hexane have the same number of electrons but pentan-1-ol is much less volatile than hexane.

Explain this difference in volatility.

(2)

.....

.....

.....

.....

.....

.....

- (c) Iodine is a solid at room temperature but on gentle heating it vaporises.

What bonds or forces are being broken?

(1)

- ☐ **A** covalent bonds
- ☐ **B** ionic bonds
- ☐ **C** London forces
- ☐ **D** permanent dipole-dipole forces



- (d) The boiling temperature of boron trichloride is 13°C while that of phosphorus trichloride is 76°C .

Explain, in terms of the shapes of the molecules, why phosphorus trichloride has a much higher boiling temperature, although both compounds have polar bonds.


Do not consider differences in London forces.

(4)

(Total for Question 2 = 8 marks)



- 3 Sodium stearate is a constituent of soap and can be made from the hydrolysis of stearin as shown.



- (a) What is the atom economy, by mass, for the production of sodium stearate in this reaction?

Substance	stearin	sodium hydroxide	sodium stearate	glycerol
Molar mass / g mol ⁻¹	890	40	306	92

(1)

- ☐ A 32.9 %
☐ B 34.4 %
☐ C 90.9 %
☐ D 98.7 %

- (b) What is the IUPAC name for glycerol?

(1)

- ☐ A 1,2,3-trihydroxypropane
☐ B propanetriol
☐ C 2,3-dihydroxypropan-1-ol
☐ D propane-1,2,3-triol

(Total for Question 3 = 2 marks)



4 Covalent bonds can break by either heterolytic fission or homolytic fission.

- (a) The reaction of butane and chlorine involves homolytic bond fission of the chlorine molecule, using ultraviolet radiation.
- (i) The energy of ultraviolet radiation can be calculated using the formula shown.

$$\text{Energy of one photon} = \frac{h \times c}{\lambda}$$

Planck constant, $h = 6.626 \times 10^{-34} \text{ J s}$

Speed of light, $c = 3.00 \times 10^8 \text{ m s}^{-1}$

Avogadro constant, $L = 6.02 \times 10^{23} \text{ mol}^{-1}$

Wavelength, λ in metres

The mean C—H bond energy is 413 kJ mol^{-1} compared to 243 kJ mol^{-1} for the Cl—Cl bond.

Show, by calculation, that ultraviolet radiation, with the wavelength 315 nm, can break the Cl—Cl bond but not the C—H bond.

(4)

- (ii) In the reaction of chlorine with butane, the propagation stage produces two different butyl radicals.

Write the structural formulae of these two radicals.

(2)

- (iii) The reaction of chlorine with butane produces 2-chlorobutane (72 %) and 1-chlorobutane (28 %).

Explain, by application of your knowledge of the stability of carbocations, why there is a higher yield of 2-chlorobutane.

(2)

.....

.....

.....

.....

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



- (b) Water molecules can undergo either heterolytic fission to form a hydroxide ion or homolytic fission to form a hydroxyl radical.

Draw dot-and-cross diagrams of these two species.

Use crosses (x) for oxygen electrons and dots (•) for hydrogen electrons.

(2)



hydroxide ion



hydroxyl radical


- (c) Which process does **not** involve heterolytic fission?

(1)

- ☐ **A** the formation of the cyanide nucleophile from KCN
- ☐ **B** the formation of propanenitrile from bromoethane
- ☐ **C** hydroxide ion attack on bromopropane in ethanol
- ☐ **D** aqueous silver nitrate in ethanol reacting with iodoethane

(Total for Question 4 = 11 marks)

- 5 The kinetics of the reactions of hydrochloric acid with metal carbonates were investigated using the apparatus shown.



The equation for the reaction is shown.



- (a) One student used 0.355 g of calcium carbonate.

Calculate the minimum volume, in cm^3 , of $0.100 \text{ mol dm}^{-3}$ hydrochloric acid that would completely react with this mass of calcium carbonate.
Give your answer to an appropriate number of significant figures.

[Calcium carbonate molar mass = 100.1 g mol^{-1}]

(4)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



(b) State why powdered metal carbonate reacts faster than large lumps.

(1)

(c) A second student used 4.19×10^{-3} mol of magnesium carbonate and suggested heating up the apparatus to speed up the rate of reaction.

- (i) If the use of heat means that the carbon dioxide gas is at a temperature of 74.0°C , then the apparatus shown would **not** give a measurable result.

Justify this comment by calculating the volume of carbon dioxide gas at this temperature using the ideal gas equation.


Assume that the pressure remains at 100 kPa throughout the experiment.

(4)


- (ii) The student suggested that the Maxwell–Boltzmann distribution of molecular energies for a reaction mixture at a particular temperature is shown in a graph.

Explain, by adding another line to the graph **and** by annotating the graph, how heating the reaction mixture results in a faster reaction rate.

(3)



- (d) A third student, in order to obtain the rate of reaction, measured the volume of gas produced at various times and used their data to plot the graph shown.



Determine the initial rate of this reaction.
You **must** show your working on the graph.
Include units in your answer.

(2)

(Total for Question 5 = 14 marks)

6 This question is about benzaldehyde and some related compounds.

* (a) Benzaldehyde ($\text{C}_6\text{H}_5\text{CHO}$) can be distinguished from phenylethanone ($\text{C}_6\text{H}_5\text{COCH}_3$) and benzoic acid ($\text{C}_6\text{H}_5\text{COOH}$) using **chemical** tests.

Describe a series of **chemical** tests which could be used to distinguish each of these compounds, giving the relevant tests and their positive results.

Do **not** use tests which give positive results for more than one of these compounds.

Detailed information of the laboratory equipment is not required.

(6)



DO NOT WRITE IN THIS AREA


DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Handwriting practice area with 20 sets of horizontal dotted lines.



(b) The reaction of 2,4-dinitrophenylhydrazine (2,4-DNPH) with a general aldehyde, RCHO, is shown.



(i) What is this type of reaction?

(1)

- ☐ **A** addition
- ☐ **B** condensation
- ☐ **C** hydrolysis
- ☐ **D** polymerisation

(ii) Draw the structure of the organic product of the reaction of 2,4-DNPH with benzaldehyde.

(1)

(Total for Question 6 = 8 marks)



7 This question is about amines.

(a) Aliphatic amines may be prepared from halogenoalkanes or from nitriles.

Assess these two methods for the preparation of butylamine.
Consider the reagents, the conditions needed, atom economy and
the type of reaction.
Include equations.

(7)

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



(b) Preparation of the aromatic amine, benzene-1,3-diamine, requires different conditions.

- (i) Write an equation for the preparation of benzene-1,3-diamine from the corresponding dinitro-compound and include necessary conditions.

(3)

Conditions




- (ii) Explain the factors affecting the basicity of benzene-1,3-diamine and propane-1,3-diamine.

Include reference to information given in the table.

Substance	pK_{b1}
benzene-1,3-diamine	8.89
propane-1,3-diamine	3.83

K_{b1} is the equilibrium constant for the first protonation reaction.

The equation is shown.



(4)

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....



- (c) Methylamine reacts with ethanoyl chloride to produce an amide.
An incomplete outline of the mechanism is shown.

Add curly arrows to show the movement of the electrons in the mechanism.


(3)



(Total for Question 7 = 17 marks)



- 8 The flavour of aniseed and liquorice is mainly due to anethole, which has the structure shown.




- (a) Give the molecular formula of anethole.

(1)

- (b) Draw the stereoisomer of anethole.

(1)


- (c) A structural isomer of anethole is estragole, which has the structure shown.



Give a reason why anethole shows stereoisomerism but estragole does **not**.

(1)

(d) What is the reagent required to convert anethole to the diol shown?



(1)

- ☐ **A** acidified potassium manganate(VII)
- ☐ **B** aqueous sodium hydroxide
- ☐ **C** ethanolic potassium hydroxide
- ☐ **D** steam with an acid catalyst

(e) The structure of anethole can be drawn with a delocalised model of the benzene ring or with the Kekulé benzene ring structure as shown.

delocalised model	Kekulé structure

- (i) Describe how the orbitals of carbon atoms overlap to form the σ -bonds and the π -bonds in the delocalised model of the benzene ring. You may include a labelled diagram in your answer.

(3)

.....

.....

.....

.....

.....



- (ii) Explain why the delocalised model of the benzene ring is the preferred representation rather than the Kekulé cyclic triene.
Refer to **one** physical and **one** thermodynamic measurement in your answer.

(4)

.....

.....

.....

.....

.....

.....

.....


.....



(f) Anethole is also present in some alcoholic drinks such as the Greek liquor, ouzo.

The anethole is responsible for the so-called 'ouzo effect'. This effect is the transparent ouzo liquor turning milky when water is added. The addition of water causes the anethole to form oily droplets which result in a stable milky emulsion.

Deduce the solubility of anethole in ethanol and in water from the 'ouzo effect'. Justify your answer with reference to the anethole structure shown, but detailed descriptions of intermolecular forces are not required.



anethole

(3)

.....

.....

.....


.....

.....

.....

.....

(g) A derivative of anethole is anisaldehyde, which has the structure shown.



What is the number of peaks in the ^{13}C NMR spectrum of anisaldehyde?

(1)

- ☐ **A** 5
- ☐ **B** 6
- ☐ **C** 7
- ☐ **D** 8

(Total for Question 8 = 15 marks)



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

BLANK PAGE



9 Halogenoalkanes are hydrolysed to form alcohols.

- (a) The rate constant for the reaction of 1-bromopropane with aqueous sodium hydroxide was determined at five temperatures. Some of the results are given in the table.

Temperature (T) / K	1/Temperature ($1/T$) / K^{-1}	Rate constant (k) / $\text{dm}^3 \text{mol}^{-1} \text{s}^{-1}$	$\ln k$
298	3.36×10^{-3}	1.36×10^{-4}	-8.9
	3.29×10^{-3}		-8.4
308	3.25×10^{-3}	3.04×10^{-4}	-8.1
318	3.14×10^{-3}	6.76×10^{-4}	-7.3
328	3.06×10^{-3}	1.23×10^{-3}	-6.7

- (i) Complete the table of data.

(2)

- (ii) Plot a graph of $\ln k$ against $1/T$ and use it to determine the activation energy, E_a , for this reaction in kJ mol^{-1} .

You must show your working on the graph and include the value of the gradient of the line and its units.

The Arrhenius equation can be expressed as

$$\ln k = -\frac{E_a}{R} \times \frac{1}{T} + \text{constant}$$


(5)



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Gradient

Activation energy



- (iii) The rate constant for the reaction of 1-bromopropane with aqueous potassium hydroxide has the units of $\text{dm}^3 \text{mol}^{-1} \text{s}^{-1}$.

Explain why it can be deduced that the reaction is second order overall. Assume that the units of the rate of reaction are $\text{mol dm}^{-3} \text{s}^{-1}$.

(2)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



(b) The rate of reaction for the hydrolysis of 2-bromo-2-methylpropane by aqueous hydroxide ions is independent of the concentration of hydroxide ions.

- (i) Draw the S_N1 reaction mechanism for this hydrolysis.
Include curly arrows and any relevant dipoles and lone pairs of electrons.

(3)

- (ii) State how your mechanism is consistent with the rate of reaction being independent of hydroxide ion concentration.

(1)

.....

.....

.....

.....

(Total for Question 9 = 13 marks)

TOTAL FOR PAPER = 90 MARKS

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

BLANK PAGE



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

BLANK PAGE



The Periodic Table of Elements

1 2 3 4 5 6 7 0 (8) (18)

1.0
H
hydrogen
1

Key

relative atomic mass
atomic symbol
name
atomic (proton) number

(1) (2)

6.9 Li lithium 3
9.0 Be beryllium 4
23.0 Na sodium 11
24.3 Mg magnesium 12

(3)

45.0 Sc scandium 21
88.9 Y yttrium 39
87.6 Sr strontium 38
137.3 Ba barium 56
132.9 Cs caesium 55

(4)

47.9 Ti titanium 22
91.2 Zr zirconium 40
178.5 Hf hafnium 72
173.3 La* lanthanum 57
[227] Ac* actinium 89

* Lanthanide series

(5)

50.9 V vanadium 23
92.9 Nb niobium 41
180.9 Ta tantalum 73
[226] Db dubnium 105

(6)

52.0 Cr chromium 24
95.9 Mo molybdenum 42
183.8 W tungsten 74
[266] Sg seaborgium 106

* Actinide series

(7)

54.9 Mn manganese 25
[98] Tc technetium 43
186.2 Re rhenium 75
[264] Bh bohrium 107

(8)

55.8 Fe iron 26
101.1 Ru ruthenium 44
190.2 Os osmium 76
[277] Hs hassium 108

* Actinide series

(9)

58.9 Co cobalt 27
102.9 Rh rhodium 45
192.2 Ir iridium 77
[268] Mt meitnerium 109

(10)

58.7 Ni nickel 28
106.4 Pd palladium 46
195.1 Pt platinum 78
[271] Ds darmstadtium 110

* Actinide series

(11)

63.5 Cu copper 29
107.9 Ag silver 47
197.0 Au gold 79
[272] Rg roentgenium 111

(12)

65.4 Zn zinc 30
112.4 Cd cadmium 48
200.6 Hg mercury 80
[272] Rg roentgenium 111

* Actinide series

(13)

10.8 B boron 5
27.0 Al aluminium 13
69.7 Ga gallium 31
114.8 In indium 49
112.4 Cd cadmium 48

(14)

12.0 C carbon 6
28.1 Si silicon 14
72.6 Ge germanium 32
118.7 Sn tin 50
207.2 Pb lead 82

* Actinide series

(15)

14.0 N nitrogen 7
31.0 P phosphorus 15
74.9 As arsenic 33
121.8 Sb antimony 51
209.0 Bi bismuth 83

(16)

16.0 O oxygen 8
32.1 S sulfur 16
79.0 Se selenium 34
127.6 Te tellurium 52
[209] Po polonium 84

* Actinide series

(17)

19.0 F fluorine 9
35.5 Cl chlorine 17
79.9 Br bromine 35
126.9 I iodine 53
[210] At astatine 85

(18)

20.2 Ne neon 10
39.9 Ar argon 18
83.8 Kr krypton 36
131.3 Xe xenon 54
[222] Rn radon 86

* Actinide series

Elements with atomic numbers 112-116 have been reported but not fully authenticated

140	Ce	cerium	58	141	Pr	praseodymium	59	144	Nd	neodymium	60	150	Sm	samarium	62	152	Eu	europium	63	157	Gd	gadolinium	64	163	Dy	dysprosium	66	165	Ho	holmium	67	167	Er	erbium	68	173	Yb	ytterbium	70	175	Lu	lutetium	71
232	Th	thorium	90	[231]	Pa	protactinium	91	238	U	uranium	92	[242]	Pu	plutonium	94	[243]	Am	americium	95	[247]	Cm	curium	96	[251]	Cf	californium	98	[254]	Es	einsteinium	99	[253]	Fm	fermium	100	[256]	Md	mendelevium	101	[257]	Lr	lawrencium	103

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

