

Paper Reference 9MA0/01
Pearson Edexcel
Level 3 GCE

Mathematics
Advanced
PAPER 1: Pure Mathematics 1

Tuesday 6 June 2023 – Afternoon

Time: 2 hours

YOU MUST HAVE
Mathematical Formulae and Statistical Tables (Green),
calculator

YOU WILL BE GIVEN
Diagram Booklet
Answer Booklet

Q72804A

Candidates may use any calculator allowed by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

INSTRUCTIONS

In the boxes on the Answer Booklet and on the Diagram Booklet, write your name, centre number and candidate number.

Answer ALL questions and ensure that your answers to parts of questions are clearly labelled.

Answer the questions in the Answer Booklet or on the separate diagrams – there may be more space than you need.

Do NOT write on this Question Paper.

You should show sufficient working to make your methods clear. Answers without working may not gain full credit.

Inexact answers should be given to three significant figures unless otherwise stated.

INFORMATION

A booklet ‘Mathematical Formulae and Statistical Tables’ is provided.

**There are 15 questions in this Question Paper.
The total mark for this paper is 100.**

**The marks for EACH question are shown in brackets
– use this as a guide as to how much time to spend on
each question.**

**There may be spare copies of some diagrams in case
you need them.**

ADVICE

**Read each question carefully before you start to answer
it.**

Try to answer every question.

Check your answers if you have time at the end.

Turn over

1. Find

$$\int \frac{x^{\frac{1}{2}}(2x-5)}{3} dx$$

writing each term in simplest form.

(Total for Question 1 is 4 marks)

2. In this question you must show all stages of your working.

Solutions relying entirely on calculator technology are not acceptable.

$$f(x) = 4x^3 + 5x^2 - 10x + 4a \quad x \in \mathbb{R}$$

where a is a positive constant.

Given $(x - a)$ is a factor of $f(x)$,

(a) show that

$$a(4a^2 + 5a - 6) = 0$$

(2 marks)

(continued on the next page)

2. continued.

(b) Hence

(i) find the value of a

(ii) use algebra to find the exact solutions of the equation

$$f(x) = 3$$

(4 marks)

(Total for Question 2 is 6 marks)

Turn over

3. Relative to a fixed origin **O**

- the point **A** has position vector $5\mathbf{i} + 3\mathbf{j} + 2\mathbf{k}$
- the point **B** has position vector $2\mathbf{i} + 4\mathbf{j} + a\mathbf{k}$

where **a** is a positive integer.

(a) Show that

$$|\vec{OA}| = \sqrt{38}$$

(1 mark)

(b) Find the smallest value of **a** for which

$$|\vec{OB}| > |\vec{OA}|$$

(2 marks)

(Total for Question 3 is 3 marks)

4. In this question you must show all stages of your working.

Solutions relying entirely on calculator technology are not acceptable.

The curve **C** has equation
 $y = f(x)$ where $x \in \mathbb{R}$

Given that

- $f'(x) = 2x + \frac{1}{2} \cos x$
- the curve has a stationary point with
 x coordinate α
- α is small

- (a) use the small angle approximation for $\cos x$ to estimate the value of α to 3 decimal places.
(3 marks)

(continued on the next page)

Turn over

4. continued.

The point

P(0, 3) lies on **C**

(b) Find the equation of the tangent to the curve
at **P**, giving your answer in the form
 $y = mx + c$, where **m** and **c** are constants to
be found.

(2 marks)

(Total for Question 4 is 5 marks)

5. Refer to the table for Question 5 in the Diagram Booklet.

A continuous curve has equation

$$y = f(x)$$

The table in the Diagram Booklet shows corresponding values of x and y for this curve, where a and b are constants.

The trapezium rule is used, with all the y values in the table, to find an approximate area under the curve between

$$x = 3 \text{ and}$$

$$x = 4$$

Given that this area is 17.59

(a) show that

$$a + 2b = 51$$

(3 marks)

(continued on the next page)

Turn over

5. continued.

Given also that the sum of all the y values in the table is 97.2

**(b) find the value of a and the value of b
(3 marks)**

(Total for Question 5 is 6 marks)

6. $a = \log_2 x$

$$b = \log_2(x + 8)$$

Express in terms of a and/or b

(a) $\log_2 \sqrt{x}$
(1 mark)

(b) $\log_2(x^2 + 8x)$
(2 marks)

(c) $\log_2\left(8 + \frac{64}{x}\right)$

Give your answer in simplest form.

(3 marks)

(Total for Question 6 is 6 marks)

Turn over

7. The function f is defined by

$$f(x) = 3 + \sqrt{x - 2} \quad x \in \mathbb{R} \quad x > 2$$

(a) State the range of f

(1 mark)

(b) Find f^{-1}

(3 marks)

The function g is defined by

$$g(x) = \frac{15}{x - 3} \quad x \in \mathbb{R} \quad x \neq 3$$

(c) Find

$gf(6)$

(2 marks)

(continued on the next page)

7. continued.

(d) Find the exact value of the constant a for which

$$f(a^2 + 2) = g(a)$$

(2 marks)

(Total for Question 7 is 8 marks)

8. Refer to the diagram for Question 8 in the Diagram Booklet.

It is NOT accurately drawn.

It shows the plan view of a stage.

The plan view shows two congruent triangles **ABO** and **GFO** joined to a sector **OCDEO** of a circle, centre **O**, where

- angle **COE** = 2.3 radians
- arc length **CDE** = 27.6 metres
- **AOG** is a straight line of length **15** metres

(a) Show that

$$\mathbf{OC = 12 \text{ metres}}$$

(2 marks)

(b) Show that the size of angle **AOB** is

$$\mathbf{0.421 \text{ radians correct to 3 decimal places.}}$$

(2 marks)

(continued on the next page)

Turn over

8. continued.

Given that the total length of the front of the stage,
BCDEF, is **35** metres,

(c) find the total area of the stage, giving your
answer to the nearest square metre.

(6 marks)

(Total for Question 8 is 10 marks)

9. The first three terms of a geometric sequence are

$$3k + 4$$

$$12 - 3k$$

$$k + 16$$

where k is a constant.

- (a) Show that k satisfies the equation

$$3k^2 - 62k + 40 = 0$$

(2 marks)

Given that the sequence converges,

- (b) (i) find the value of k , giving a reason for your answer,

(ii) find the value of S_{∞}

(5 marks)

(Total for Question 9 is 7 marks)

10. A circle C has equation

$$x^2 + y^2 + 6kx - 2ky + 7 = 0$$

where k is a constant.

(a) Find in terms of k ,

(i) the coordinates of the centre of C

(ii) the radius of C

(3 marks)

The line with equation

$y = 2x - 1$ intersects C at 2 distinct points.

(b) Find the range of possible values of k

(6 marks)

(Total for Question 10 is 9 marks)

11. Refer to the diagram for Question 11 in the Diagram Booklet.

The value, V pounds, of a mobile phone, t months after it was bought, is modelled by

$$V = ab^t$$

where a and b are constants.

The diagram shows the linear relationship between $\log_{10} V$ and t

The line passes through the points $(0, 3)$ and $(10, 2.79)$ as shown.

Using these points,

- (a) find the initial value of the phone,
(2 marks)**

(continued on the next page)

11. continued.

- (b) find a complete equation for V in terms of t ,
giving the exact value of a and giving the
value of b to 3 significant figures.
(3 marks)**

**Exactly 2 years after it was bought, the value of the
phone was £320**

- (c) Use this information to evaluate the reliability
of the model.
(2 marks)**

(Total for Question 11 is 7 marks)

12. $y = \sin x$

where x is measured in radians.

Use differentiation from first principles to show that

$$\frac{dy}{dx} = \cos x$$

You may

- use without proof the formula for $\sin(A \pm B)$
- assume that as

$$h \rightarrow 0,$$

$$\frac{\sin h}{h} \rightarrow 1 \text{ and}$$

$$\frac{\cos h - 1}{h} \rightarrow 0$$

(Total for Question 12 is 5 marks)

Turn over

13. On a roller coaster ride, passengers travel in carriages around a track.

On the ride, carriages complete multiple circuits of the track such that

- the maximum vertical height of a carriage above the ground is **60** metres
- a carriage starts a circuit at a vertical height of **2** metres above the ground
- the ground is horizontal

The vertical height, **H** metres, of a carriage above the ground, **t** seconds after the carriage starts the first circuit, is modelled by the equation

$$H = a - b(t - 20)^2$$

where **a** and **b** are positive constants.

(continued on the next page)

13. continued.

(a) Find a complete equation for the model.

(3 marks)

(b) Use the model to determine the height of the carriage above the ground when $t = 40$

(1 mark)

In an alternative model, the vertical height, H metres, of a carriage above the ground, t seconds after the carriage starts the first circuit, is given by

$$H = 29 \cos(9t + \alpha)^\circ + \beta \quad 0 \leq \alpha < 360^\circ$$

where α and β are constants.

(continued on the next page)

13. continued.

(c) Find a complete equation for the alternative model.

(2 marks)

Given that the carriage moves continuously for 2 minutes,

(d) give a reason why the alternative model would be more appropriate.

(1 mark)

(Total for Question 13 is 7 marks)

14. Prove, using algebra, that

$$(n + 1)^3 - n^3$$

is odd for all $n \in \mathbb{N}$

(Total for Question 14 is 4 marks)

15. A curve has equation
 $y = f(x)$, where

$$f(x) = \frac{7xe^x}{\sqrt{e^{3x} - 2}} \quad x > \ln \sqrt[3]{2}$$

- (a) Show that

$$f'(x) = \frac{7e^x(e^{3x}(2-x) + Ax + B)}{2(e^{3x} - 2)^{\frac{3}{2}}}$$

where **A** and **B** are constants to be found.

(5 marks)

- (b) Hence show that the **x** coordinates of the turning points of the curve are solutions of the equation

$$x = \frac{2e^{3x} - 4}{e^{3x} + 4}$$

(2 marks)

(continued on the next page)

15. continued.

The equation

$$x = \frac{2e^{3x} - 4}{e^{3x} + 4} \text{ has two positive roots } \alpha \text{ and } \beta$$

where $\beta > \alpha$

A student uses the iteration formula

$$x_{n+1} = \frac{2e^{3x_n} - 4}{e^{3x_n} + 4}$$

in an attempt to find approximations for α and β

Refer to the diagram for Question 15(c) in the Diagram Booklet.

It shows a plot of part of the curve with equation

$$y = \frac{2e^{3x} - 4}{e^{3x} + 4} \text{ and part of the line with equation}$$

$$y = x$$

(continued on the next page)

Turn over

15. continued.

Using the diagram in the Diagram Booklet,

- (c) draw a staircase diagram to show that the iteration formula starting with $x_1 = 1$ can be used to find an approximation for β**
(1 mark)

Use the iteration formula with $x_1 = 1$, to find, to 3 decimal places,

- (d) (i) the value of x_2**

- (ii) the value of β**

(3 marks)

(continued on the next page)

15. continued.

Using a suitable interval and a suitable function that should be stated

(e) show that

$\alpha = 0.432$ to 3 decimal places.

(2 marks)

(Total for Question 15 is 13 marks)

TOTAL FOR PAPER IS 100 MARKS

END OF PAPER
