

**Physics**

**Advanced**

**Paper 1: Advanced Physics I**

<b>Total Marks</b>
--------------------

**Monday 18 May 2020 – Afternoon**

**Time: 1 hour 45 minutes plus your additional time allowance**

**In the boxes below, write your name, centre number and candidate number.**

<b>Surname</b>					
<b>Other names</b>					
<b>Centre Number</b>					
<b>Candidate Number</b>					

## **YOU MUST HAVE**

**Scientific calculator, ruler, protractor**

## **YOU WILL BE GIVEN**

**Diagram Booklet, Data, formulae and relationships Booklet**

## **INSTRUCTIONS**

**Answer ALL questions.**

**Answer the questions in the spaces provided – there may be more space than you need.**

## **INFORMATION**

**The total mark for this paper is 90.**

**The marks for EACH question are shown in brackets – use this as a guide as to how much time to spend on each question.**

**You may use a scientific calculator.**

**In questions marked with an ASTERISK (\*), marks will be awarded for your ability to structure your answer logically, showing how the points that you make are related or follow on from each other where appropriate.**

**A data, formulae and relationships booklet is provided.**

**ADVICE**

**Read each question carefully before you start to answer it.**

**Try to answer every question.**

**Check your answers if you have time at the end.**

**You are advised to show your working in calculations including units where appropriate.**

**Answer ALL questions.**

**All multiple choice questions must be answered with a cross ☒ in the box for the correct answer from A to D. If you change your mind about an answer, put a line through the box ~~☒~~ and then mark your new answer with a cross ☒.**

**1 Which of the following is an example of a scalar quantity?**

☐ **A displacement**

☐ **B energy**

☐ **C momentum**

☐ **D velocity**

**(TOTAL FOR QUESTION 1 = 1 MARK)**

---

- 2 Which row of the table summarises the mass and charge of an antineutron?

	Mass / u	Charge / e
<input type="checkbox"/> A	0	0
<input type="checkbox"/> B	0	-1
<input type="checkbox"/> C	1	0
<input type="checkbox"/> D	1	+1

(TOTAL FOR QUESTION 2 = 1 MARK)

---

- 3 The distance between a proton and an electron is  $r$ .  
The electrostatic force is  $F$ .

The distance between the proton and electron is doubled.

Which of the following is equal to the electrostatic force at this separation?

☐ A  $2F$

☐ B  $\frac{F}{2}$

☐ C  $\frac{F}{3}$

☐ D  $\frac{F}{4}$

(TOTAL FOR QUESTION 3 = 1 MARK)

---

4 Which of the following is a base SI unit?

- ☐ A ampere
- ☐ B coulomb
- ☐ C joule
- ☐ D newton

(TOTAL FOR QUESTION 4 = 1 MARK)

---

5 A high-energy proton can interact with a photon to produce two particles.

Which of the following could be the two particles produced?

- ☐ A  $n + \pi^0$
- ☐ B  $n + \pi^+$
- ☐ C  $\pi^0 + \pi^+$
- ☐ D  $\pi^- + \pi^+$


(TOTAL FOR QUESTION 5 = 1 MARK)

---


- 6 Look at the graph for Question 6 in the Diagram Booklet. It shows how an electric potential  $V$  varies with distance  $x$ .

Which of the following shows the corresponding variation of electric field strength  $E$  with  $x$ ?


☐ A




☐ B



☐ C



☐ D



(TOTAL FOR QUESTION 6 = 1 MARK)

---



- 7 The intensity of light incident on a light dependent resistor (LDR) can vary both its electrical resistance  $R$  and the number of charge carriers per unit volume  $n$ . The light intensity on an LDR is increased.

Which row of the table describes the effect on  $R$  and  $n$ ?

	$R$	$n$
<input type="checkbox"/> A	decreases	decreases
<input type="checkbox"/> B	decreases	increases
<input type="checkbox"/> C	increases	decreases
<input type="checkbox"/> D	increases	increases

(TOTAL FOR QUESTION 7 = 1 MARK)

---

- 8 A proton has a mass of  $1.67 \times 10^{-27}$  kg.

Which of the following shows the conversion of this mass to  $\text{GeV}/c^2$  ?

☐ A 
$$\frac{1.67 \times 10^{-27} \times 1.60 \times 10^{-10}}{(3.00 \times 10^8)^2}$$

☐ B 
$$\frac{1.67 \times 10^{-27} \times 1.60 \times 10^{-19}}{(3.00 \times 10^8)^2}$$

☐ C 
$$\frac{1.67 \times 10^{-27} \times (3.00 \times 10^8)^2}{1.60 \times 10^{-10}}$$

☐ D 
$$\frac{1.67 \times 10^{-27}}{1.60 \times 10^{-10} \times (3.00 \times 10^8)^2}$$

(TOTAL FOR QUESTION 8 = 1 MARK)

---

- 9 The blade of a lawnmower rotates at a speed of 50 revolutions per second.

Which of the following is the angular speed of the blade in  $\text{rads s}^{-1}$ ?

☐ A 7.96

☐ B 15.9

☐ C 157

☐ D 314

(TOTAL FOR QUESTION 9 = 1 MARK)

---

- 10 Two gliders, X and Y, are placed on an air track.  
Look at the diagram for Question 10 in the Diagram Booklet. The gliders are pushed towards each other.

The gliders collide and continue to move after the collision.

Which row of the table could show the velocities of X and Y, in  $\text{ms}^{-1}$ , after the collision?

	X	Y
<input type="checkbox"/> A	-1.0	0.5
<input type="checkbox"/> B	-1.0	-0.5
<input type="checkbox"/> C	-2.0	-1.0
<input type="checkbox"/> D	-2.0	2.0

(TOTAL FOR QUESTION 10 = 1 MARK)

---

**11 A uniform paving slab is to be used as a garden step.**

- (a) State what is meant by the centre of gravity of an extended body. (1 mark)**

---

---

---

- (b) Look at the diagram for Question 11(b) in the Diagram Booklet. The paving slab has a weight of 310 N and a length of 90 cm and will be supported at two points, P and Q. The distance between P and Q will be 75 cm.**

**This might be unsafe because a person who places all their weight at the end of the slab might tip the slab.**

**A person of mass 70 kg stands at the end of the slab.**

**Deduce whether the slab will tip. (4 marks)**

---

---

**11 continued.**

---

---

---

---

---

---

---

---

---

---

---

---

**(TOTAL FOR QUESTION 11 = 5 MARKS)**

---

- 12** Analogue ammeters were used before digital meters became widely available. On page 7 in the Diagram Booklet the analogue ammeter will measure a maximum current of  $1.0\text{ mA}$  and has a resistance of  $18\Omega$ .

The analogue ammeter can be adapted to measure a larger current by adding a resistor, known as a shunt, in parallel with the ammeter. On page 7 in the Diagram Booklet the arrangement is shown. The analogue ammeter is represented by the  $18\Omega$  resistor.

The maximum current through the  $18\Omega$  resistor remains as  $1.0\text{ mA}$ .

- (a) Show that the shunt would need to have a resistance of about  $0.01\Omega$  to adapt this ammeter to read up to a maximum current of  $2.0\text{ A}$ . (3 marks)

---

---

---

---

---

---

**12 continued.**

---

---

---

**(continued on the next page)**



**12 continued.**

- (b) A shunt of this resistance was usually made from Manganin wire.**

**Calculate the length of Manganin wire of radius 0.95 mm required to make this shunt. (3 marks)**

**resistivity of Manganin =  $4.55 \times 10^{-7} \Omega \text{m}$**

---

---

---

---

---

---

---

---

---

---

**Length = \_\_\_\_\_**

**(TOTAL FOR QUESTION 12 = 6 MARKS)**

**Turn over**

**13 Look at the image for Question 13 in the Diagram Booklet. It shows a model racing car set. The curved parts of the track are semicircular. The car makes electrical contact with the track using metal brushes underneath the car.**

- (a) There is a maximum speed for the car to stay on the curved part of the track. Explain why the car will slip off the curved part of the track if the car exceeds the maximum speed. (3 marks)**

---

---

---

---

---

---

---

---

---

---

**(continued on the next page)**

**13 continued.**

- (b) The following measurements are made for a car starting at rest on a straight piece of track.**

**distance travelled = 1.2 m**

**time taken = 0.77 s**

- (i) Show that the final velocity of the car is about  $3 \text{ m s}^{-1}$ .**

**Assume the acceleration is constant.  
(2 marks)**

---

---

---

---

---

**(continued on the next page)**

13 continued.

- (ii) The final velocity calculated in (b)(i) is the maximum velocity before the car slips off the track.

Calculate the maximum horizontal force between the curved part of the track and the car. (2 marks)

mass of car = 0.050 kg

radius of curved part of track = 0.042 m

---

---

---

---

---

---

Maximum horizontal force = \_\_\_\_\_

(continued on the next page)

**13 continued.**

- (c) The cars are controlled separately and so can be raced, with one car on the inner lane and the other on the outer lane. The cars are identical. Each car is raced at its highest speed for that lane.**

**Explain why the outcome of the race is difficult to predict. (3 marks)**

---

---

---

---

---

---

---

---

---

---

**(TOTAL FOR QUESTION 13 = 10 MARKS)**

---

- 14 On page 9 in the Diagram Booklet the diagram shows a model used to demonstrate alpha particle scattering. A ball bearing is set rolling on a wooden track. The track is positioned so that the ball bearing rolls onto a metal sheet with a curved surface known as a 'hill'.

On page 9 in the Diagram Booklet the diagram shows a vertical cross-section through the hill. The surface is curved so that the height of a point  $h$  on the curved surface is inversely proportional to the distance  $r$  from the centre of the hill.

- (a) Explain why the hill is suitable as a model for the electric field surrounding the nucleus of an atom.  
(3 marks)

---

---

---

---

---

---

---

(continued on the next page)

Turn over

**14 continued.**

- (b) Look at the diagram for Question 14(b) in the Diagram Booklet. It shows a plan view of the arrangement.**

**The wooden track is moved to different positions and the ball bearing is released.**

**Describe the results of the alpha particle scattering experiment and how these can be demonstrated by moving the wooden track to different positions.  
(4 marks)**

---

---

---

---

---

---

---

---

---

**(continued on the next page)**

**Turn over**

**14 continued.**

---

---

---

---

**(TOTAL FOR QUESTION 14 = 7 MARKS)**

---



- 15** The properties of capacitors make them useful in timing circuits.

Look at the circuit for Question 15 in the Diagram Booklet. It is used to provide an input Y to an integrated circuit.

- (a)** Initially the capacitor is uncharged. The switch  $S_1$  is closed.

Look at the grid for Question 15(a) in the Diagram Booklet. Sketch a graph to show how the potential at point Y varies with time. (3 marks)

---

---

---

**(continued on the next page)**

**15 continued.**

**(b) When the potential at Y is 8.0 V, the switch  $S_2$  is closed.**

**(i) Calculate the time taken for the potential at Y to decrease to 2.0 V. (3 marks)**

---

---

---

---

---

---

---

**Time taken = \_\_\_\_\_**

**(continued on the next page)**

15 continued.

- (ii) Calculate the energy stored on the capacitor when the potential at Y is 2.0 V. (2 marks)

---

---

---

---

---

---

Energy stored = \_\_\_\_\_

(continued on the next page)

**15 continued.**

- (c) When the potential at Y is 2.0 V, the switch  $S_2$  is opened.**

**Calculate the power dissipated by the resistance  $R_1$  when the potential at Y is 2.0 V.  
(3 marks)**

---

---

---

---

---

---

---

---

---

---

**Power dissipated = \_\_\_\_\_**

**(TOTAL FOR QUESTION 15 = 11 MARKS)**

**Turn over**

**16 Scientists have been studying a type of jumping spider that can jump up to six times its body length.**

**(a) Look at the picture for Question 16(a) in the Diagram Booklet. The scientists photographed a spider at 0.02 s intervals, during a jump. The picture is taken from the photograph and has been enlarged.**

**(i) Deduce whether the images show that the motion in the x-direction is independent of the motion in the y-direction. You should take measurements using the cross marking the centre of gravity of the spider. (4 marks)**

---

---

---

---

---

---

---

**(continued on the next page)**

**16 continued.**

---

---

---

---

---

**(continued on the next page)**

**16 continued.**

- (ii) Show that the initial velocity of the spider at the start of the jump is about  $1 \text{ m s}^{-1}$ . You should take measurements using the cross marking the centre of gravity of the spider. (5 marks)**

---

---

---

---

---

---

---

---

---

---

---

**(continued on the next page)**

**Turn over**

**16 continued.**

---

---

---

---

---

---

**(continued on the next page)**



**16 continued.**

- (iii) The spider achieves this jump by extending its two back legs by 3.0 mm.**

**Calculate the average force the spider exerts in each leg to achieve the jump. (3 marks)**

**mass of spider = 150 mg**

---

---

---

---

---

---

**Average force = \_\_\_\_\_**

**(continued on the next page)**

**16 continued.**

- (b) Look at the image for Question 16(b) in the Diagram Booklet. Just as the spider starts the jump, it fixes a silk thread to the platform. It is thought that the thread acts as a safety line in case the spider falls.**

**A student makes the comment:**

**‘If the silk thread can withstand a tension equal to the weight of the spider then this safety system should work.’**

**Deduce whether this statement is correct. (2 marks)**

---

---

---

---

---

---

---

---

---

**(TOTAL FOR QUESTION 16 = 14 MARKS)**

**Turn over**

**17 Hybrid electric vehicles (HEV) use the same device both as a generator to charge the car battery and as an electric motor to support the propulsion system. Look at the simplified diagram of the device for Question 17(a) in the Diagram Booklet. The coil can rotate freely around the axis.**

**\*(a) Describe how the device can be used as both a generator and an electric motor. (6 marks)**

---

---

---

---

---

---

---

---

---

---

**(continued on the next page)**

**Turn over**

17 continued.

---

---

---

---

---

---

---

---

---

(continued on the next page)

17 continued.

- (b) Look at the circuit diagram for Question 17(b) in the Diagram Booklet. It shows a car battery connected to an electric motor for a HEV. The battery has an electromotive force (e.m.f.) 180 V and internal resistance  $0.036\Omega$ .

The motor has a maximum power of 88 kW.

- (i) Show that the current  $I$  drawn by the electric motor when operating at this power would be given by the equation (3 marks)

$$0.036I^2 - 180I + 88\,000 = 0$$

---

---

---

---

---

---

---

---

(continued on the next page)

Turn over

**17 continued.**

---

---

**(continued on the next page)**

**17 continued.**

- (ii) Solving the equation above produces an answer of  $I = 550\text{ A}$ . At maximum power, the car can accelerate from rest to sixty miles per hour in under 7 s.

The maximum charge capacity of the battery within this HEV is 6.1 amp-hour.

Deduce whether the battery could maintain this current for up to 7 s. (2 marks)

---

---

---

---

---

---

---

**(TOTAL FOR QUESTION 17 = 11 MARKS)**

---

**18 A negatively charged pion decays into a muon and an antineutrino. Look at the diagram for Question 18 in the Diagram Booklet. It shows tracks in a particle detector formed in such an event.**

**(a) Deduce whether the antineutrino is charged, giving two reasons for your decision. (2 marks)**

---

---

---

---

---

---

**(b) Write a particle equation to represent this decay. (1 mark)**

---

---

**(continued on the next page)**



**18 continued.**

- (c) According to the standard model, the pion and muon are classified within two different groups of particles.**

**State which group each particle belongs to and describe the two groups. (4 marks)**

---

---

---

---

---

---

---

---

---

---

---

---

18 continued.

- (d) The momentum of the pion just before it decays is  $9.1 \times 10^{-20} \text{ N s}$ .

Determine the magnetic flux density of the magnetic field which acts in the detector and state its direction. (4 marks)

Scale of diagram 1 cm represents 10 cm

pion charge =  $-1.6 \times 10^{-19} \text{ C}$

---

---

---

---

---

---

---

---

---

(continued on the next page)

Turn over

18 continued.

---

---

---

---

Magnetic flux density = \_\_\_\_\_

Direction of magnetic field = \_\_\_\_\_

(continued on the next page)

**18 continued.**

- (e) Use a vector diagram to determine the momentum of the antineutrino.**

**The initial momentum of the muon is  $1.59 \times 10^{-19} \text{ N s}$ .  
(5 marks)**

**(continued on the next page)**

**Turn over**

18 continued.

Momentum of antineutrino = \_\_\_\_\_

Direction of antineutrino = \_\_\_\_\_

(TOTAL FOR QUESTION 18 = 16 MARKS)

---

---

TOTAL FOR PAPER = 90 MARKS

END