

Physics  
Advanced  
PAPER 2: Advanced Physics II

Total Marks
-------------

Thursday 6 June 2024 – Morning

Time: 1 hour 45 minutes

In the boxes below, write your name, centre number and candidate number.

Surname					
Other names					
Centre Number					
Candidate Number					

## **YOU MUST HAVE**

**Scientific calculator and ruler**

**Data, Formulae and Relationships Booklet (enclosed)**

## **YOU WILL BE GIVEN**

**Diagram Booklet**

## **INSTRUCTIONS**

**Answer ALL questions.**

**Answer the questions in the spaces provided in this Question Paper or in the separate Diagram Booklet – there may be more space than you need.**

## **INFORMATION**

**The total mark for this paper is 90.**

**The marks for EACH question are shown in brackets – use this as a guide as to how much time to spend on each question.**

**In the question marked with an ASTERISK (\*), marks will be awarded for your ability to structure your answer logically, showing how the points that you make are related or follow on from each other where appropriate.**

**There may be spare copies of some diagrams.**

## **ADVICE**

**Read each question carefully before you start to answer it.**

**Try to answer every question.**

**Check your answers if you have time at the end.**

**You are advised to show your working in calculations, including units where appropriate.**

---

**Answer ALL questions.**

**All multiple choice questions must be answered with a cross in the box ☒ for the correct answer from A to D. If you change your mind about an answer, put a line through the box ☒ and then mark your new answer with a cross ☒.**

- 1 Look at the diagram for Question 1 in the Diagram Booklet. A Hertzsprung–Russell diagram for stars in our galaxy is shown. V, W, X, Y and Z are positions on the diagram.**

**Which of the following gives a sequence of positions of the Sun during its lifetime?**

☐ **A V X Z**

☐ **B X W Y**

☐ **C X W Z**

☐ **D Z X V**

**(Total for Question 1 = 1 mark)**

---

2 A satellite of mass  $m$  orbits the Earth with speed  $V$ .

What is the speed of a satellite of mass  $2m$  orbiting at the same distance from the Earth?

☐ A  $\frac{V}{2}$

☐ B  $V$

☐ C  $V\sqrt{2}$

☐ D  $2V$

(Total for Question 2 = 1 mark)

---

- 3 The isotope  $^{221}_{87}\text{Fr}$  undergoes a series of alpha and beta decays before forming  $^{205}_{81}\text{Tl}$ .

Which row of the table shows the number of alpha and beta particles emitted during the decays?

	Alpha	Beta
<input type="checkbox"/> A	3	2
<input type="checkbox"/> B	3	6
<input type="checkbox"/> C	4	2
<input type="checkbox"/> D	4	6

(Total for Question 3 = 1 mark)

---

- 4 In an electron diffraction experiment, electrons are accelerated by a potential difference between a hot filament and an anode. The electron beam produced strikes a sample of crystalline material causing a diffraction pattern.**

**Which of the following would cause the angle of diffraction to increase?**

- ☐ **A decreasing the distance between the filament and the anode**
- ☐ **B decreasing the potential difference**
- ☐ **C increasing the filament temperature**
- ☐ **D using a crystalline material with a larger lattice spacing**

**(Total for Question 4 = 1 mark)**

---

- 5 A particle **P** has charge and mass. The particle causes an electric field and a gravitational field.

Which of the following statements is correct, at a distance **r** from **P**?

☐ A gravitational field strength is proportional to  $\frac{1}{r}$

☐ B electric field strength is proportional to  $\frac{1}{r}$

☐ C gravitational potential is always positive

☐ D electric potential is proportional to  $\frac{1}{r}$

(Total for Question 5 = 1 mark)

---



- 6 Look at the graph for Question 6 in the Diagram Booklet. It shows how binding energy per nucleon varies with nucleon number for atomic nuclei. The arrows **W**, **X**, **y** and **Z** represent changes in binding energy per nucleon and nucleon number.

Fission and fusion of nuclei may result in the release of energy.

Which row of the table gives the arrows representing energy release by fission and fusion?

	Fission	Fusion
<input type="checkbox"/> A	<b>x</b>	<b>w</b>
<input type="checkbox"/> B	<b>x</b>	<b>y</b>
<input type="checkbox"/> C	<b>z</b>	<b>w</b>
<input type="checkbox"/> D	<b>z</b>	<b>y</b>

(Total for Question 6 = 1 mark)

---

- 7 Look at the diagram for Question 7 in the Diagram Booklet. It shows a ray of light passing from medium **X** to medium **Y** to medium **Z**. The speed of light in medium **X** is  $v_X$ , the speed of light in medium **Y** is  $v_Y$  and the speed of light in medium **Z** is  $v_Z$ .

Which of the following shows the speeds in order of increasing magnitude?

☐ A  $v_Y$   $v_X$   $v_Z$

☐ B  $v_Z$   $v_X$   $v_Y$

☐ C  $v_X$   $v_Y$   $v_Z$

☐ D  $v_Y$   $v_Z$   $v_X$

(Total for Question 7 = 1 mark)

---

- 8 A weight of **2 N** is suspended from a spring of spring constant **50 N m<sup>-1</sup>**

What is the elastic strain energy stored by the spring in joules?

☐ A  $\frac{1}{2} \times 2 \times 50$

☐ B  $\frac{1}{2} \times 2 \times 50^2$

☐ C  $\frac{1}{2} \times \frac{2}{50}$


☐ D  $\frac{1}{2} \times \frac{2^2}{50}$

(Total for Question 8 = 1 mark)

---

- 9 A mass is suspended from a spring and displaced vertically. The mass performs simple harmonic motion.

The graph shows how the acceleration of the mass varies with time over one oscillation.



Look at the graphs for Question 9 in the Diagram Booklet. Which of these graphs shows how the velocity  $V$  of the mass varies with time  $t$  over the same time interval?

☐ A

☐ B

☐ C

☐ D

(Total for Question 9 = 1 mark)

---

- 10 A wire of length  $L$  and diameter  $d$  is fixed at one end. A force  $F$  is applied to the wire causing an extension  $e$ .

A second wire, made of the same material, of length  $2L$  and diameter  $\frac{d}{2}$ , is fixed in the same way. A force  $2F$  is applied to this wire.

What is the extension of the second wire?

☐ A  $\frac{e}{4}$

☐ B  $e$

☐ C  $8e$

☐ D  $16e$

(Total for Question 10 = 1 mark)

---

- 11 A student shone green light from a laser through a diffraction grating, producing a diffraction pattern on a screen. The student determined the angle of the third order maximum.

Calculate the wavelength of the green light.

angle of third order maximum =  $73.3^\circ$

grating spacing =  $1.67 \times 10^{-6} \text{ m}$

---

---

---

---

---

---

Wavelength = \_\_\_\_\_

(Total for Question 11 = 2 marks)

---

- 12 Study of the photoelectric effect leads to observations that cannot be explained by the wave theory of light.**

**When ultraviolet light is shone on a charged zinc plate, electrons are released. When visible light is shone on the zinc plate, electrons are not released.**

- (a) (i) State what is meant by threshold frequency.  
(1 mark)**

---

---

---

- (ii) Explain why the observation of a threshold frequency is NOT consistent with the wave theory of light.  
(2 marks)**

**Answer space continues on the next page.**

---

---

---

---

**12(a)(ii) continued.**

---

---

**(b) The photoelectric effect can be explained using the idea of photons.**

**State what is meant by the term photon.  
(1 mark)**

---

---

---

**(Total for Question 12 = 4 marks)**

---



**13 Aldebaran is a red giant star.**

**For a black body radiator of Aldebaran's surface temperature, the wavelength  $\lambda_{\text{max}}$  at which the intensity is maximum is  $7.43 \times 10^{-7} \text{ m}$ .**

**Determine the luminosity of Aldebaran.  
(4 marks)**

**radius of Aldebaran =  $3.14 \times 10^{10} \text{ m}$**

**Answer space continues on the next page.**

---

---

---

---

---

---

---

---

---

---

**13 continued.**

**radius of Aldebaran =  $3.14 \times 10^{10}$  m**

---

---

---

**Luminosity = \_\_\_\_\_**

**(Total for Question 13 = 4 marks)**

---

**14 A student used two converging thin lenses in combination to form an image of an object.**

- (a) Show that the power of the combination of lenses was about 20 D.  
(3 marks)**

**focal length of lens 1 = 10 cm**

**focal length of lens 2 = 15 cm**

---

---

---

---

---

---

---

---

---

---

**(continued on the next page)**

**14 continued.**

**(b) The object was 4 cm in front of the combination of lenses.**

**Calculate the image distance.  
(2 marks)**

---

---

---

---

---

---

**Image distance = \_\_\_\_\_**

**(continued on the next page)**

**14 continued.**

**(c) The object height was 1.4 cm.**

**Calculate the image height.  
(3 marks)**

---

---

---

---

---

---

---

---

**Image height = \_\_\_\_\_**

**(continued on the next page)**

**14 continued.**

**(d) The object is magnified.**

**State TWO other properties of the image.  
(1 mark)**

---

---

---

**(Total for Question 14 = 9 marks)**

---

- 15 Look at Diagram 1 for Question 15 in the Diagram Booklet. A student wound a piece of string around the head of an electric toothbrush. The student attached a small mass to the other end of the string, as shown in Diagram 1.**

**Look at Diagram 2 for Question 15 in the Diagram Booklet. The toothbrush was switched on and the head started to vibrate. The student rotated the toothbrush slowly to unwind the string. At a particular length, large vibrations were observed on the string. The string formed a loop, as shown in Diagram 2.**

- (a) Explain how the vibrations of the toothbrush head caused the loop to form.  
(5 marks)**

**Answer space continues on the next page.**

---

---

---

---

---

---

15(a) continued.

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

(continued on the next page)



**15 continued.**

- (b) The student continued to unwind the string and the loop disappeared. When the length of the unwound string was twice that shown in Diagram 2 two loops were seen. Three loops were seen when the unwound length was three times that shown in Diagram 2 and so on.**

**Determine the frequency of vibration of the toothbrush head.**

**(5 marks)**

**unwound length of string with 4 loops = 0.69 m**

**mass on string = 0.010 kg**

**mass per unit length of string =  $9.1 \times 10^{-4} \text{ kg m}^{-1}$**

**Answer space continues on the next page.**

---

---

---

---

---

---

---

**Turn over**

15(b) continued.

unwound length of string with 4 loops = **0.69 m**

mass on string = **0.010 kg**

mass per unit length of string =  **$9.1 \times 10^{-4} \text{ kg m}^{-1}$**

---

---

---

---

---

---

---

---

---

---

Frequency = \_\_\_\_\_

(Total for Question 15 = 10 marks)

---

- 16 On the International Space Station (**ISS**), astronauts measure their mass once a month using a Body Mass Measurement Device (**BMMD**).

Look at the diagram for Question 16 in the Diagram Booklet. The **BMMD** is constructed from a large spring attached to the floor of the **ISS**, with a platform and handles attached to the spring. The spring is compressed and the astronaut puts his body onto the platform and holds onto the handles, as shown.

The spring is released and the astronaut and platform oscillate with simple harmonic motion.

An astronaut used the **BMMD**. The frequency of oscillation was **0.34 Hz**.

- (a) Determine the mass of the astronaut.  
(4 marks)

spring constant =  **$350 \text{ N m}^{-1}$**

mass of platform =  **$5.7 \text{ kg}$**

Answer space continues on the next page.

---

---

---

16(a) continued.

spring constant =  $350 \text{ N m}^{-1}$

mass of platform =  $5.7 \text{ kg}$

---

---

---

---

---

---

---

---

---

---

Mass of astronaut = \_\_\_\_\_

(continued on the next page)

**16 continued.**

**(b) The distance between the upper and lower points of the first oscillation is  $0.29\text{ m}$ .**

**(i) Calculate the magnitude of the maximum acceleration of the astronaut.  
(4 marks)**

**Answer space continues on the next page.**

---

---

---

---

---

---

---

---

---

---

**16(b)(i) continued.**

---

---

**Magnitude of  
maximum acceleration = \_\_\_\_\_**

- (ii) Calculate the speed of the astronaut  $3.5\text{ s}$   
after the start of the oscillations.  
(2 marks)**

---

---

---

---

---

---

**Speed = \_\_\_\_\_**

**(Total for Question 16 = 10 marks)**

---

- 17 (a) Look at the diagram for Question 17(a) in the Diagram Booklet. A hot air balloon consists of a fabric envelope, heaters and a basket, as shown.**

**When the balloon is set up, the envelope is partly filled with air at  $20^{\circ}\text{C}$ . The air is then heated to  $120^{\circ}\text{C}$  and expands to fill the envelope and becomes less dense.**

**The air pressure inside the envelope is always equal to the air pressure outside the envelope because the envelope is open at the bottom.**

**The balloon takes off when the upthrust is more than the total weight of the balloon, the air in the envelope and the passengers.**

**Deduce whether the balloon can take off.  
(6 marks)**

**volume of air at  $120^{\circ}\text{C}$  in inflated envelope =  $2800\text{ m}^3$**

**density of air at  $20^{\circ}\text{C}$  =  $1.2\text{ kg m}^{-3}$**

**mass of balloon =  $380\text{ kg}$**

**mass of passengers =  $340\text{ kg}$**

**upthrust when the envelope is full =  $33\,000\text{ N}$**

**Answer space continues on the next 3 pages.**

17(a) continued.

volume of air at  $120^{\circ}\text{C}$  in inflated envelope =  $2800\text{ m}^3$

density of air at  $20^{\circ}\text{C}$  =  $1.2\text{ kg m}^{-3}$

mass of balloon =  $380\text{ kg}$

mass of passengers =  $340\text{ kg}$

upthrust when the envelope is full =  $33\,000\text{ N}$



**17(a) continued.**

**volume of air at  $120^{\circ}\text{C}$  in inflated envelope =  $2800\text{ m}^3$**

**density of air at  $20^{\circ}\text{C}$  =  $1.2\text{ kg m}^{-3}$**

**mass of balloon =  $380\text{ kg}$**

**mass of passengers =  $340\text{ kg}$**

**upthrust when the envelope is full =  $33\,000\text{ N}$**

---

---

---

---

---

---

---

---

---

---

---

---

**17(a) continued.**

**volume of air at  $120^{\circ}\text{C}$  in inflated envelope =  $2800\text{ m}^3$**

**density of air at  $20^{\circ}\text{C}$  =  $1.2\text{ kg m}^{-3}$**

**mass of balloon =  $380\text{ kg}$**

**mass of passengers =  $340\text{ kg}$**

**upthrust when the envelope is full =  $33\,000\text{ N}$**

---

---

---

---

**(continued on the next page)**

**17 continued.**

- (b) (i) State one assumption of the kinetic theory of gases.  
(1 mark)**

---

---

---

**(continued on the next page)**

17(b) continued.

(ii) Derive an equation to show that, for a gas at temperature  $T$ , the mean kinetic energy of the molecules =  $\frac{3}{2} kT$

(2 marks)

---

---

---

---

---

---

(continued on the next page)

17(b) continued.

- (iii) Calculate the root-mean-square speed of nitrogen molecules at a temperature of **120 °C**.  
(3 marks)

mass of nitrogen molecule = **28 u**

---

---

---

---

---

---

---

---

---

---

Root-mean-square speed = \_\_\_\_\_

(Total for Question 17 = 12 marks)

---

**18 In 1864, William Huggins and William Miller used dark lines in the spectrum of the Sun to identify elements in the Sun's atmosphere.**

**\*(a) Explain how gases in the Sun's atmosphere cause dark lines in the spectrum corresponding to different elements.**

**(6 marks)**

**Answer space continues on the next page.**

---

---

---

---

---

---

---

---

---

---

**18(a) continued.**

---

---

---

---

---

---

---

---

**(continued on the next page)**

**18 continued.**

- (b) Look at the diagram for Question 18(b) in the Diagram Booklet. It shows some energy levels of a hydrogen atom.**

**The absorption spectrum for hydrogen includes a set of lines that all derive from transitions involving the  $n = 2$  energy level. One of these lines is known as the hydrogen-alpha line.**

**Deduce the transition involved in the formation of the hydrogen-alpha line.  
(4 marks)**

**wavelength of hydrogen-alpha line = 656.46 nm**

**Answer space continues on the next page.**

---

---

---

---

---

---



**18(b) continued.**

**wavelength of hydrogen-alpha line = 656.46 nm**

---

---

---

---

---

---

**(continued on the next page)**

**18 continued.**

- (c) In 1868, William Huggins analysed light from the star Sirius A. The wavelength of the hydrogen-alpha line for light from Sirius A was slightly different from the hydrogen-alpha line observed from a source in a laboratory.**

**Huggins suggested that this difference could be explained using the Doppler effect and could be used to determine the speed and direction of the star's motion relative to the Earth.**

- (i) Assess Huggins's suggestion.  
(3 marks)**

**Answer space continues on the next page.**

---

---

---

---

---

---

---

---

**Turn over**

**18(c)(i) continued.**

---

---

**(continued on the next page)**

**18(c) continued.**

- (ii) Sirius A has a component of velocity away from the Earth of  $5.5 \text{ km s}^{-1}$**

**The wavelength of the hydrogen-alpha line observed from a source in the laboratory is  $656.46 \text{ nm}$ .**

**Calculate the wavelength of the hydrogen-alpha line as seen in the spectrum of Sirius A.  
(2 marks)**

**Answer space continues on the next page.**

---

---

---

---

---

---

---

---

**18(c)(ii) continued.**

---

---

**Wavelength =** \_\_\_\_\_

**(Total for Question 18 = 15 marks)**

---

**19 Nuclear decay is described as being spontaneous and random.**

**(a) (i) State what is meant by spontaneous and random in this context.  
(2 marks)**

---

---

---

---

---

---

**(continued on the next page)**

**19(a) continued.**

- (ii) Explain why the decay constant of an isotope can be determined even though nuclear decay is random.  
(2 marks)**

---

---

---

---

---

---

**(continued on the next page)**

**19 continued.**

- (b) A radioactive source used in a school laboratory emits alpha and beta radiation.**

**Describe how the percentage of the activity due to beta radiation may be determined using a Geiger–Müller tube and ratemeter.**

**(4 marks)**

**Answer space continues on the next page.**

---

---

---

---

---

---

---

---

---

---

**Turn over**



**19(b) continued.**

---

---

**(continued on the next page)**

**19 continued.**

**(c) Americium-241 is used in schools as a source of alpha radiation. A pure americium-241 source was bought 34 years ago by a school.**

- (i) Determine the percentage of the initial activity that would be expected today for the americium-241 source.  
(3 marks)**

**half-life of americium-241 = 432 years**

**Answer space continues on the next page.**

---

---

---

---

---

---

---

---

19(c)(i) continued.

half-life of americium-241 = 432 years

---

Expected percentage  
of initial activity = \_\_\_\_\_

(continued on the next page)

**19(c) continued.**

- (ii) The decay products of americium are unstable and undergo a series of further decays.**

**Look at the table for Question 19(c)(ii) in the Diagram Booklet. It shows the first three decays in this sequence.**

**A student states, “Protactinium-233 emits beta particles when it decays, so by now the americium-241 source bought 34 years ago will be emitting a significant amount of beta radiation.”**

**Discuss the student’s statement.  
(3 marks)**

**Answer space continues on the next page.**

---

---

---

---

---

---

**19(c)(ii) continued.**

---

---

---

**(Total for Question 19 = 14 marks)**

---

---

**TOTAL FOR PAPER = 90 MARKS**  
**END OF PAPER**