

Mark Scheme (Results) Summer 2010

IGCSE

IGCSE Chemistry (4335) Paper 2H

Edexcel is one of the leading examining and awarding bodies in the UK and throughout the world. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers.

Through a network of UK and overseas offices, Edexcel's centres receive the support they need to help them deliver their education and training programmes to learners.

For further information, please call our GCE line on 0844 576 0025, our GCSE team on 0844 576 0027, or visit our website at www.edexcel.com.

If you have any subject specific questions about the content of this Mark Scheme that require the help of a subject specialist, you may find our **Ask The Expert** email service helpful.

Ask The Expert can be accessed online at the following link:

<http://www.edexcel.com/Aboutus/contact-us/>

Alternately, you can speak directly to a subject specialist at Edexcel on our dedicated Science telephone line: 0844 576 0037

(If you are calling from outside the UK please dial + 44 1204 770 696 and state that you would like to speak to the **Science** subject specialist).

Summer 2010

Publications Code UG024205

All the material in this publication is copyright

© Edexcel Ltd 2010

SECTION A

Question			Mark	Acceptable answers	Notes	Total	
1	a	i	M1	bubbles / fizzing / effervescence / metal gets smaller / white trail	Ignore metal dissolves / gas produced Reject all answers in a(ii)	1	
		ii	M1	melts / forms a ball / darts / moves (on surface) / floats	Ignore reference to flames Reject all answers in a(i)	1	
	b	i	M1	calcium hydroxide		1	
		ii	M1	NaOH		1	
	c		M1	hydrogen / H ₂	Ignore H	1	
				M2	(squeaky) pop with burning splint / burns with a (squeaky) pop	Accept other words such as explosion / lighted spill or taper Reject glowing splint Ignore references to air/splint extinguished No CONSEQ from wrong gas	1
	d	i	M1	blue / purple	Ignore qualifiers such as light / dark / bright	1	
			M2	OH ⁻ / hydroxide	Ignore hydroxyl	1	
			ii	M1	yellow / orange	Ignore qualifiers such as light / dark / golden / bright Reject all other colours	1

Question		Mark	Acceptable answers	Notes	Total		
2	a	M1	hydrogen peroxide		1		
		M2	manganese(IV) oxide / manganese dioxide		1		
	b	M1	(gas) syringe		1		
	c	M1	catalyst / to speed up the reaction / lower activation energy		1		
	d	i	M1	(s) for both PbS and PbSO ₄		1	
			M2	(aq) for H ₂ O ₂ and (l) for H ₂ O		1	
		ii	M1	PbS / lead sulphide / sulphide ion / S ²⁻ / sulphur in lead sulphide	Ignore oxidation numbers if given	1	
			M2	gains oxygen/O/O ₂ increase in oxidation state	only award if M1 correct or sulphur ignore loss of electrons	1	
	e	i	M1	S + O ₂ → SO ₂	Ignore state symbols Accept S ₂ or S ₈	1	
			ii	M1	acidic / (forms) H ⁺ (ions) / sulphurous acid / sulphuric(IV) acid	Reject sulphuric acid / sulphuric(VI) acid	1
			iii	M1	orange		1
			M2	green	Accept blue-green	1	

Question		Mark	Acceptable answers	Notes	Total
3	a	M1	electron transfer	All marks can be scored from suitably annotated diagrams Award 0/3 if any reference to sharing electrons Ignore covalent M3 dependent on M2	1
		M2	from magnesium/Mg to chlorine/Cl		1
		M3	Mg loses two electrons and (each) Cl gains one electron		1
	b	M1	magnesium / Mg		1
		M2	loss of electrons / increase in oxidation state	Ignore number of electrons M2 independent of M1	1
	c	M1	+ and - ions / oppositely charged ions / Mg^{2+} and Cl^{-}	Need idea of + and - charge	1
		M2	strong (electrostatic) attractions (within lattice)	accept strong (ionic) bonds reject covalent bonds / molecular attraction	1
		M3	<u>lot of</u> energy needed to overcome attractions / break bonds / separate ions	Do not accept “loosening bonds” Ignore “hard to break”	1
				any mention of “intermolecular” or “intramolecular” loses M1 and M2 So “strong intermolecular forces need lots of energy to overcome” scores M3	

Question			Mark	Acceptable answers	Notes	Total
4	a	i	M1	fractional distillation / fractionation		1
		ii	M1	crude oil heated	M1 given even if describe laboratory process. Only M1 possible if describe lab process or mention cracking/breaking bonds	1
			M2	(vapour) passed into column/tower	If crude oil heated in fractionating column, then give only 1 mark for M1 and M2	1
			M3	fractions collected at different heights		1
			M4	correct reference to boiling point / molecular size / temperature gradient/hot at bottom cooler at top	Do not award if specified temperature gradient is wrong way round	1
					All marks can be gained from a suitable diagram	
	b	i	M1	bitumen		1
		ii	M1	gasoline		1
		iii	M1	bitumen		1
		iv	M1	refinery gases	Accept answers in either order	1
			M2	fuel oil	Accept naphtha in place of either	1
	c		M1	oxygen	Ignore air	1
			M2	carbon dioxide	Accept answers in either order	1
			M3	water	Accept steam in place of water	1
					All marks in c are independent	
					Ignore heat	

	d	i	M1	C_nH_{2n+2}	Accept other letters/symbols such as x accept $C_nH_{2(n+1)}$	1
		ii	M1	same/similar chemical properties / same functional group	reject trend in chemical properties	
			M2	gradation in physical properties / gradation in specified physical property (eg boiling point)	reject same/similar physical properties	
			M3	neighbouring members differ by CH_2	Accept any two for 1 mark each Ignore references to general formula and references to saturation/unsaturation/specific functional group	2

SECTION A TOTAL: 45 MARKS

SECTION B

Question		Mark	Acceptable answers	Notes	Total
5	a		M1 number of protons in an atom	Do not award mark if no mention of atom/nucleus Ignore reference to electrons unless clearly added to number of protons	1
	b	i	M1 isotope(s)		1
		ii	M1 38		1
			M2 18		1
			M3 18		1
			M4 22		1
		iii	M1 full outer energy level/shell / complete octet / no need to gain or lose electrons / eight electrons in outer energy level/shell / 2.8.8	Ignore reference to stability/ionisation energy	1
	c	i	M1 (atoms of isotope 65 are) 30.9 %		1
			M2 $(63 \times 0.691) + (65 \times 0.309)$	CONSEQ on incorrect percentage in M1	1
			M3 63.6	Correct final answer scores 3 marks Award 2 marks for 63.62 / 63.618 CONSEQ on incorrect percentage in M1 ignore units	1
		ii	M1 same electronic configuration/number of (outer) electrons	ignore references to same number of protons	1

Question		Mark	Acceptable answers	Notes	Total
6	a	M1	$\text{MgCO}_3 \rightarrow \text{MgO} + \text{CO}_2$	reagent = 1	1
		M2		both products = 1	1
				Award 1 mark for all formulae correct in an unbalanced equation ignore state symbols	
	b	i	M1	magnesium chloride/nitrate/sulphate/other soluble magnesium salt	1
			M2	sodium/potassium/ammonium carbonate / other soluble carbonate	1
		ii	M1	correct balanced equation such as $\text{MgCl}_2 + \text{Na}_2\text{CO}_3 \rightarrow \text{MgCO}_3 + 2\text{NaCl}$	2
			M2		
		iii	M1	filter / centrifuge and decant	1
			M2	Wash (residue/solid) with water	M2 and M3 dependent on an attempt at M1(eg “sieving”, “decant”)
			M3	dry by warming gently / leave (in warm place) to dry / uses filter/absorbent paper / dry in (warm) oven / place in dessicator	1
				Points must be in correct order to score all marks	

Question		Mark	Acceptable answers	Notes	Total
7	a	M1	bromine (water)	Reject bromide, but mark M2 and M3 as if bromine accept KMnO_4	1
		M2	(stays) yellow / orange / brown / no change/ no reaction	Reject red Purple if KMnO_4	1
		M3	(becomes) colourless / decolourised	Ignore clear ignore discoloured Decolourised if acidified KMnO_4 brown if neutral KMnO_4 green if alkaline KMnO_4 if only KMnO_4 allow any of above three accept 1,2-dibromopropane (if bromine) or propan(e)-1,2-diol (if KMnO_4)	1
	b	M1	alkene(s)	Accept olefins	1
	c	M1	$\begin{array}{cc} \text{H} & \text{CH}_3 \\ & \\ \text{---C---} & \text{C---} \\ & \\ \text{H} & \text{H} \end{array}$	M1 for correct structure (ignore continuation bonds)	1
		M2	$\begin{array}{cc} \text{---C---} & \text{C---} \\ & \\ \text{H} & \text{H} \end{array}$	M2 for continuation bonds	1
				M2 dependent on M1 Ignore brackets and subscript letters Award 0 marks if double bond shown	
	d	M1	poly(propene) / polypropene / polypropylene		1

Question			Mark	Acceptable answers	Notes	Total
8	a	i	M1	$\text{Zn} + \text{CuSO}_4 \rightarrow \text{Cu} + \text{ZnSO}_4$	M1 for reagents	1
			M2	$/ \text{Zn} + \text{Cu}^{2+} \rightarrow \text{Cu} + \text{Zn}^{2+}$	M2 for products	1
					Ignore state symbols Award 1 for all formulae correct in unbalanced equation	
		ii	M1	(copper is) less reactive (than zinc)/lower (in reactivity series than zinc) /	Accept "copper forms ions less easily" Accept reverse argument for zinc Reject answers that compare reactivity of ions.	1
		iii	M1	(red-)brown/pink solid/ppt/coating(on zinc)	Accept copper in place of colour	1
			M2	solution becomes colourless/ paler		1
	b	i	M1	sacrificial (protection/anode)	Ignore galvanising	1
		ii	M1	zinc is more reactive than iron/steel/hull / higher in reactivity series than iron/steel/hull	Accept reverse argument for iron/steel Accept "they" for zinc blocks	1
			M2	zinc reacts (with air/water) instead of/ before/ in preference to iron/steel/hull /prevents iron from losing electrons/zinc makes Fe^{2+} gain electrons	reject zinc rusts reject references to a protective coating of zinc or zinc oxide If have zinc sacrificing itself here, can award mark for (i) if not contradictory to (i)	1
		iii	M1	copper less reactive than iron/steel/hull / lower in reactivity series than iron/steel/hull / copper does not react with air/water / copper makes iron corrode more / copper makes iron lose electrons	They = copper blocks Accept converse argument	1

Question		Mark	Acceptable answers	Notes	Total	
9	a	M1	$\text{PCl}_5 + 4\text{H}_2\text{O} \rightarrow \text{H}_3\text{PO}_4 + 5\text{HCl}$	M1 for all four formulae correct	1	
		M2		M2 for balancing	1	
				M2 dependent on M1 Allow multiples and fractions		
	b	M1	(starts) green		1	
		M2	(turns) red/pink	accept orange if only 1 colour given and not clear whether start or end, then do not award mark.	1	
		M3	(becomes) acidic / acid / H^+ (ions) (formed)	Accept $\text{pH} < 7$	1	
	c	i	M1	two atoms linked by shared pair of electrons	Atoms do not have to be labelled H and Cl,	1
			M2	six more electrons in Cl and no more electrons in H	Ignore inner electrons in Cl M2 dependent on M1 do not award M2 if atoms are wrongly identified Accept any suitable symbol(s) for electrons	1
		ii	M1	Shared electrons/ electron pair		1
			M2	attracts/pulls nucleus/nuclei / protons (in nuclei)	ignore neutrons	1
	d		M1	weak forces of attraction		1
			M2	between molecules/intermolecular	Idea of covalent bonds breaking = 0 Intramolecular bonds are covalent therefore breaking them scores 0 Weak intermolecular bonds = 2	1
	e	i	M1	$0.150 \times 12.3/1000$	Ignore units	1
			M2	$0.001845 / 0.00185$	Award max 1 mark wrong conversion or no conversion of volume to dm^3 correct answer alone scores 2	1
		ii	M1	$(0.001845 \times 3 =) 0.005535$	CONSEQ on e(i)/M2	1
		iii	M1	$\frac{0.005535}{0.025}$	CONSEQ on e(ii) Award 1 mark for 0.0002214 (not divided vol by 1000)	1
			M2	0.2214	Final answer must be to 2 or more sig fig Ignore units	1

					Correct final answer scores 2 marks	
					If correct answer of 0.2214 obtained by failing to convert cm^3 to dm^3 correctly in both ei and eiii, then max 1 in both ei and eiii	

Question		Mark	Acceptable answers	Notes	Total
10	a		M1 phosphoric acid / H ₃ PO ₄	Ignore dilute / concentrated	1
			M2 high temperature / 250-350 °C / high pressure / 50-100 atmospheres	Ignore heat If range of T or P given, must be fully within accepted range If give both a temperature and a pressure , then do not apply list principle	1
	b	i	M1 3186	Accept 1075	1
		ii	M1 3231	Accept 1120	1
				Ignore units and signs	
		iii	M1 -45	CONSEQ on bi - bii Sign must be shown if negative Note - if (i) correct and (ii) 2883, then conseq answer in (iii) is (+)303.	1
	c		M1 $\begin{array}{c} \text{H H H} \\ \\ \text{H-C-C=C} \\ \quad \\ \text{H} \quad \text{H} \end{array}$	Accept CH ₃ CHCH ₂ as minimum	1

Question			Mark	Acceptable answers	Notes	Total
11	a	i	M1	0.1(00)		1
		ii	M1	(M_r of CuCO_3 =) 123.5 / 124		1
			M2	$0.1(00) \times 123.5$ or $124 = 12.35$ or 12.4		1
					CONSEQ on ai Correct final answer scores 2 marks If final answer wrong, and M1 not awarded, award M2 for showing multiplication of any number by a(i)	
		iii	M1	$0.1(00) \times 24$ / $0.1(00) \times 24000$	CONSEQ on answer to ai	1
			M2	$= 2.4 \text{ (dm}^3\text{)}$ / $= 2400 \text{ cm}^3$	Correct final answer with units scores 2 marks If no units stated assume dm^3	1
	b		M1	(light) blue precipitate	Accept ppt / solid in place of precipitate Reject dark/deep/royal	1
			M2	dark(er) blue solution	Accept deep / royal in place of dark	1
			M3	$[\text{Cu}(\text{NH}_3)_4(\text{H}_2\text{O})_2]^{2+}$	Square brackets not needed NH_3 and H_2O can be in reverse order	1

Question		Mark	Acceptable answers	Notes	Total
12	a		M1 lattice/regular arrangement/array of positive ions	If positive AND negative ions do not award M1	1
			M2 sea of/delocalised/mobile electrons	Accept free electrons	1
			M3 electrons move	If ions move then do not award M3	1
	b	i	M1 steel/iron reacts with chlorine /forms iron chloride /oxidises	ignore references to corrodes /wears away	1
		ii	M1 titanium / Ti		1
	c	i	M1 $100\,000 \times 2 \times 60$		1
			M2 12 000 000	Correct answer alone scores 2 Award 1 mark for answer of 200 000 Ignore units	1
		ii	M1 $\frac{12\,000\,000}{96\,000} = 125$		1
			M2 $\frac{125}{2} = 62.5 / 63$	CONSEQ on answer from ci correct answer scores 2 if fail to convert coulombs to Faradays, can score M2 only (6 000 000)	1

SECTION B TOTAL: 75 MARKS

PAPER TOTAL: 120 MARKS

Further copies of this publication are available from
International Regional Offices at www.edexcel.com/international

For more information on Edexcel qualifications, please visit www.edexcel.com
Alternatively, you can contact Customer Services at www.edexcel.com/asktheexpert or on + 44 1204 770 696

Edexcel Limited. Registered in England and Wales no.4496750
Registered Office: One90 High Holborn, London, WC1V 7BH